

THE HYSTERO-ONTOGENY OF *LONSDALEIA*  
MCCOY AND *THYSANOPHYLLUM ORIENTALE*  
THOMSON

by R. K. JULL

**ABSTRACT.** Increase in thirteen specimens of *Lonsdaleia* and two of *Thysanophyllum orientale* from Great Britain was studied by means of closely spaced acetate peel sections. With the exception of one example of intermural increase, only lateral increase was observed. *Lonsdaleia* is characterized during early development by having a long cardinal septum, the axial end of which forms the median lamella of the axial structure. Septal insertion is at first of the zaphrentid mode but subsequently becomes variable, depending on the development of minor septa. Most daughter corallites are radially orientated to the parent, but some are tangential. Interseptal dissepiments are typically formed first, followed by their modification to lonsdaleoid dissepiments.

Development in *Thysanophyllum orientale*, type species of *Thysanophyllum*, resembles that in *Lonsdaleia* in the nature of the cardinal septum and dissepimental development. These characters, together with the presence in a few adult corallites of *T. orientale* of a weakly developed axial structure suggest that *Thysanophyllum* is related to the Lonsdaleiidae rather than the Lithostrotionidae as has been suggested by some workers.

SOME of the taxonomic problems resulting from the morphologic variation of rugose corals have not been satisfactorily solved by the standard technique of examining adult characters alone. Such difficulties exist as regards the family affinities of *Thysanophyllum* Nicholson and Thomson, a widely distributed genus of Carboniferous and Permian age which lacks an axial structure and has short septa and lonsdaleoid dissepiments (Pl. 102, figs. 5a-b). Some workers (Lecompte 1952, Hill 1956, Fedorowski 1965) regard this genus as being a diphymorphic lonsdaleoid and thus place it with the Lonsdaleiidae. Others, however (Minato 1955, Easton 1960, Fontaine 1961, Soshkina and Dobrolyubova 1962), assign it to the Lithostrotionidae partly since corallites of *Lithostrotion* Fleming may lack an axial structure and possess lonsdaleoid dissepiments, thus closely resembling the characters of *Thysanophyllum*.

A solution to this problem is attempted here by a study of closely spaced acetate peel sections of the characters of increase in thirteen specimens of *Lonsdaleia* and two of *Thysanophyllum orientale*, type species of *Thysanophyllum*. These specimens, which are listed in text-fig. 1, are all from Viséan beds in Great Britain. From a comparison of the hystero-ontogeny of *Thysanophyllum orientale* with that in *Lithostrotion* (see Jull 1965) and *Lonsdaleia*, typical genera respectively of the Lithostrotionidae and Lonsdaleiidae, it is concluded that *Thysanophyllum* is probably related to the Lonsdaleiidae.

Observations of hystero-corallite development range from the start of increase up to the late neanic stage and sometimes the ephebic stage. Crushing in three coralla has destroyed the early stages of development in some or all of the corallites studied.

The terminology applied by Jull (1965) to increase in *Lithostrotion* is used herein. Abbreviations associated with catalogue numbers are as follows: BM—British Museum (Natural History), London; RSM—Royal Scottish Museum, Edinburgh; UQ—Department of Geology and Mineralogy, University of Queensland, Brisbane.

*Acknowledgements.* I am grateful to Dr. J. W. Wells (Cornell University) for his criticism of the manuscript, and to Professor D. Hill (University of Queensland) for her help during the early stages of this [Palaeontology, Vol. 10, Part 4, 1967, pp. 617-28, pls. 100-102.]

study and for permitting use of specimens in her care. Dr. C. D. Waterston (Royal Scottish Museum), Mr. I. F. Sime (Edinburgh), Dr. C. T. Scrutton (British Museum (Natural History)), and Mr. G. P. McDonagh (Phillips Petroleum, Brisbane) all kindly contributed specimens for the study. I also wish to acknowledge the co-operation of the University of Queensland where this study was initiated, and Cornell University where it was completed.

Name	Number	Locality	Zonal age	Corallites studied	Remarks
<i>Lonsdaleia duplicata duplicata</i> (Martin)	BM R1366	Corwen, North Wales	D <sub>2</sub> -D <sub>3</sub>	4	
	RSM 1878.11.267	Hafod, near Corwen, North Wales	D <sub>2</sub> -D <sub>3</sub>	3	early stages crushed
<i>L. floriformis floriformis</i> (Martin)	UQ F11035	Wrekin, Shropshire	D <sub>2</sub>	1	diphymorphic corallites
	UQ F2961	Pen-y-ghent, Yorkshire	D <sub>2</sub> -D <sub>3</sub>	3	
	RSM 1966.6.1	Coldstone Quarries, Greenhow, Pately Bridge, Yorkshire	D <sub>2</sub>	8	
	BM R17160	Coalbrookdale, Shropshire	D <sub>2</sub>	5	
	BM R24850	Longstone Edge, north of Bakewell, Derbyshire	D <sub>2</sub>	2	diphymorphic corallites
<i>L. floriformis crassiconus</i> McCoy	UQ F5683	Clifton, near Bristol	D <sub>2</sub>	8	
	RSM 1966.6.2	Gathorn Plain, Orton, Westmorland	D <sub>2</sub>	14	
	BM R16995	Avon Gorge, Bristol	D <sub>2</sub>	17	topotype
	BM R16997	Avon Gorge, Bristol	D <sub>2</sub>	13	topotype
<i>L. floriformis near laticlavata</i> Smith	BM R36865	Oswestry, Shropshire	D <sub>2</sub> -D <sub>3</sub>	3	
<i>L. caledonia</i> Smith	RSM 1966.6.3	Coalburn, Lesmahagow, Lanarkshire, Scotland	D <sub>2</sub>	3	early stages crushed
<i>Thysanophyllum orientale</i> Thomson	UQ F46767	Aberlady Bay, East Lothian, Scotland	Coral Zone 3	6	topotype
	RSM 1966.6.4	West Kirkton Quarry, Bathgate, West Lothian, Scotland	Coral Zone 3	7	early stages crushed in 3

TEXT-FIG. 1. Specimens of *Lonsdaleia* McCoy and *Thysanophyllum orientale* Thomson studied by means of closely spaced acetate peel sections.

#### INCREASE IN *LONSDALEIA*

Studies of increase in Lower Carboniferous representatives of *Lonsdaleia* have been made by thin section examination of British material by Smith (1916) and on material from the Russian Platform by Dobrolyubova (1958). Fedorowski (1965) described development in two Lower Permian species of *Lonsdaleia* from Hornsund, Vestspitsbergen, using serial acetate peel sections. These studies include a large number of species but there are nevertheless significant aspects of the hystero-ontogeny requiring clarification. These are mainly: (1) the orientation of the axial plane of the daughter corallite with respect to the axis of the parent; (2) the manner of development of lonsdaleoid dissepiments; (3) the pattern of septal insertion; and (4) the nature of the wall dividing

the daughter from the parent corallite. These characters, together with present and past knowledge of other characters, are dealt with in turn in the following discussion. Species are discussed collectively since there generally seems to be little fundamental difference between their hystero-ontogenies. As text-fig. 1 indicates, most of the observations are made on *L. floriformis floriformis* and *L. floriformis crassiconus*. Complete details of increase were obtained from only one specimen each of *L. floriformis* near *laticlavia* and *L. duplicata duplicata*, and the early characters of development are unknown in *L. caledonia*.

*Mode of increase.* This is almost invariably lateral in nature with the daughter corallite arising in the lonsdaleoid dissepimentarium of the parent in no preferred position with respect to the axial plane of the parent. Commonly as many as five daughter corallites may develop nearly simultaneously from the same parent (Pl. 102, fig. 2), especially in the proximal and lateral parts of the corallum. The rate of corallite development in cerioid species varies greatly but usually about 20–30 mm. of growth occurs between the start of development and the achievement of the ephebic stage. Development in the fasciculate *L. duplicata duplicata* is considerably slower, extending over some 40 mm. or more of distal growth with about 35 mm. of this occurring during late neanic development.


Rarely intermural increase (*sensu* Jull 1965, p. 206) also occurs whereby a youthful corallite is formed between adjacent corallites and lacks a single (obvious) parent. One example of this type of increase was observed in *L. floriformis floriformis* (text-fig. 2; Pl. 102, figs. 4 *a, b*), and Dobrolyubova (1958, p. 79–81, pl. 9, fig. 1; text-figs. 14–16) described and illustrated this type of increase in *L. rossica rossica* Stuckenbergh from Lower Namurian C<sub>1</sub><sup>st</sup> beds on the Russian Platform.

*Dividing wall.* In all examples of increase observed, the wall which divides the daughter from the parent corallite is similar to the wall bounding adult corallites. It is formed either by thickening at the peripheral ends of parent septa (Pl. 100, figs. 2 *b–f*), a character similar to that noted by Fedorowski (1965, p. 121), or by apparent modification of one or more lonsdaleoid dissepiments of the parent (corallite D3, Pl. 101, figs. 1 *a–c*). When corallites of *L. duplicata duplicata* (text-fig. 4, 1 *a–j*) become discontinuous, this wall is split rather than progressively disappearing as does the partition which divides daughter from parent in most fasciculate species of *Lithostrotion*.

*Septal insertion.* In the specimens studied, septa are inserted independently of the parent septa. Usually the cardinal septum is the first to appear and the other primary septa, the counter septum and two alar septa, may not all be present when metasepta first start to appear, especially if the dividing wall is late in forming. Insertion is of a zaphrentid nature in all quadrants (Pl. 100, figs. 2 *f–j*) and this may either continue up to the ephebic stage or minor septa may appear during development and insertion alters in all quadrants to the cyathaxonid mode (Pl. 100, figs. 2 *k–m*). The development of minor septa is discussed below.

In contrast to septal formation in Lower Carboniferous forms, Fedorowski (1965, p. 121) noted that the earliest septa in daughter corallites of the two Lower Permian species that he studied are formed from the detached ends of parent septa. In both species parent septa extend to the wall in the zone of increase preceding the appearance

of the daughter corallite whereas in the Lower Carboniferous forms, this zone of increase is usually free of parent septa.



TEXT-FIG. 2. Intermural increase in *Lonsdaleia floriformis floriformis* (Martin), UQ F11035. Numbers below figures are cumulative distances of distal growth in mm. Thick lines are intercorallite walls and septa; thin lines are dissepiments and tabulae. In A, the youthful corallite is formed in lonsdaleoid dissepiments and is only separated from adult corallites lying to the upper-right and lower-left by lonsdaleoid dissepiments. Neither of these adult corallites could be established as the parent. The union of axial septa in C-H is uncommon in *Lonsdaleia*. Developmental stages, of which the late neanic is the last illustrated (Q), merge into one another.  $\times 4$ .

*Axial septa and axial structure.* During early development, the cardinal septum is almost invariably the longest septum and the counter and alar septa are typically indistinguishable from metasepta. The median lamella of the axial structure is formed from the axial

end of the cardinal septum, and during the progressive appearance of tabellae and other lamellae in the axial structure, the cardinal septum may or may not remain attached to the median lamella. During the ephelic stage and later part of the late neanic stage the two are usually detached.

Dobrolyubova (1958, p. 27) did not trace septal insertion in the species which she examined but suggested that the longer septum was the cardinal septum. Smith (1916, pp. 230, 232), on the other hand, considered the longer septum to be the counter septum but he likewise did not trace septal insertion. In some of the youthful corallites which he illustrated (*ibid.*, pl. 17, figs. 5–23), insertion appears to be taking place beside this septum, indicating that it is perhaps the cardinal septum. De Groot (1963, p. 80) remarked that an elongate cardinal septum is characteristic of lonsdaleoid genera, but Fedorowski (1965) did not note any long septum during development in his lonsdaleoid species.

Name	Number	Radial	Tan- gential	Indeter.	Total
<i>L. duplicata</i> <i>duplicata</i> (Martin)	BM R1366	2	1(?)	1	4
	RSM 1878.11.267	—	—	3	3
<i>L. floriformis</i> <i>floriformis</i> (Martin)	UQ F2961	3	—	—	3
	RSM 1966.6.1	6	1	1	8
	BM R17160	1	2	2	5
	BM R24850	2	—	—	2
<i>L. floriformis</i> <i>crassiconus</i> McCoy	UQ F5683	7	1	—	8
	RSM 1966.6.2	10	3	1	14
	BM R16995	14	1	2	17
	BM R16997	11	—	2	13
<i>L. floriformis</i> near <i>laticlavata</i> Smith	BM R36865	1	1	1	3
<i>L. caledonia</i> Smith	RSM 1966.6.3	—	—	3	3

TEXT-FIG. 3. The number of corallites with radial and tangential orientation in the specimens studied. Orientation is based on the position of the axial plane of the daughter corallite with respect to the axis of the parent. Indeterminate corallites are those in which the early stages are not preserved, usually through crushing, or the position of the axial plane could not be definitely established.

*Orientation of the daughter corallite.* Text-fig. 3 shows that the majority of corallites studied are radially orientated with the cardinal septum positioned on the outer side, furthest from the axis of the parent. Most of these corallites are orientated at less than 30° from a true radial orientation (Pl. 100; corallites D2 and D3, Pl. 101), a few are up to 45° away from this position (corallite D1, Pl. 101) and those that are tangentially orientated lie at from 60° to 90° away from a radial position (Pl. 102, figs. 1*a-c*; text-fig. 4, 2*a-e*). Preference to radial orientation is exemplified by those specimens in which up to five daughter corallites, all radially orientated, are ranged simultaneously around the periphery of the parent (Pl. 102, fig. 2). With one exception in which two corallites develop together, all tangentially orientated daughter corallites observed arise singly.

In many of Dobrolyubova's (1958) illustrations, the axial plane of the daughter corallite appears to be radial to the parent, but in *L. arctica* Gorsky (ibid., text-fig. 3), *L. subtilis* Dobrolyubova (ibid., text-fig. 19) and *L. floriformis floriformis* (ibid., text-fig. 12) radial orientation appears to be lacking. Possibly this may be due to the axial plane of the daughter corallite rotating during development, as I have observed in the Viséan *Aphrophyllum hallense* Smith, but this effect has not been observed in the specimens of *Lonsdaleia* here studied.

Radial orientation of the daughter to parent corallite is certainly the most common condition in *Lonsdaleia* but non-radial orientation is too common to be an abnormal occurrence. Perhaps orientation of the daughter to parent corallite is a factor of individual variation, with radial orientation being the most common tendency.

*Dissepiments.* The first one or two rows of dissepiments in the daughter corallite are typically formed first as interseptal dissepiments when ten or more major septa have been inserted. These are subsequently altered to lonsdaleoid dissepiments by the withdrawal of septa from the wall. In a few cases, however, lonsdaleoid dissepiments of the parent corallite are incorporated as such into the morphology of the daughter (text-fig. 4, 1*a-j*).

*Minor septa.* Minor septa in *Lonsdaleia* are variably developed in different specimens. In some examples of increase, especially those in which major septa are mainly confined

---

EXPLANATION OF PLATE 100

Numbers below Figs. 2*a-m* are cumulative distances of distal growth in mm. C = cardinal septum; K = counter septum; A = alar septum.

Figs. 1, 2. *Lonsdaleia floriformis floriformis* (Martin). 1, UQ F2961, transverse section of normal and diphymorphic corallites,  $\times 1$ . 2*a-m*, lateral increase in the same specimen; *a-f*, hystero-brephic stage; *g-l*, early neanic stage, showing the appearance of the counter septum in *g* and of minor septa in *k*; *m*, late neanic stage, also showing another youthful corallite with a long cardinal septum on the right,  $\times 4$ .

EXPLANATION OF PLATE 101

Numbers below figures are cumulative distances of distal growth in mm. D = daughter corallite; P = parent corallite; C = cardinal septum; K = counter septum.

Figs. 1*a-h*. *Lonsdaleia floriformis crassiconus* McCoy, UQ F5683. Lateral increase in three corallites, with daughter corallites D1, D2, and D3 arising from parent corallites P1, P2, and P3 respectively. Note that the axial plane of D3 is nearly radially disposed with respect to the axis of its parent, while in D1 and D2, it is approximately 20 to 30 degrees away from a radial disposition. Minor septa in D3 appear in *f-h*, and interseptal dissepiments are formed in *d*, followed by their alteration to lonsdaleoid dissepiments in *e-g*,  $\times 4$ .

EXPLANATION OF PLATE 102

Numbers below Figs. 1*a-c* are cumulative distances of distal growth in mm. D = daughter corallite; P = parent corallite; C = cardinal septum; K = counter septum.


Figs. 1-3. *Lonsdaleia floriformis crassiconus* McCoy. 1*a-c*, UQ F5683, laterally arising corallite with its axial plane orientated tangential to the axis of the parent; *a, b*, early neanic stage; *c*, late neanic stage,  $\times 4$ . 2, BM R16997, five radially orientated daughter corallites arising from the same parent,  $\times 4$ . 3, transverse view of the same specimen as in Figs. 1*a-c*,  $\times 1$ .

Figs. 4*a, b*. *Lonsdaleia floriformis floriformis* (Martin). UQ F11035, transverse and longitudinal sections of diphymorphic corallites,  $\times 1$ .

Figs. 5*a, b*. *Thysanophyllum orientale* Thomson. Topotype, UQ F46767, transverse and longitudinal sections,  $\times 1$ .

---

to the tabular area, such as in *L. duplicata duplicata* and *L. floriformis floriformis*, minor septa are absent or very short throughout hystero-ontogeny, so that insertion is entirely zaphrentid in nature. In others, however, minor septa appear as early as simultaneously with the development of the first dissepiments, in which case insertion alters to the cyathaxonid mode.




TEXT-FIG. 4. Numbers below figures are cumulative distances of distal growth in mm. C = cardinal septum. 1 a-j, lateral increase in *Lonsdaleia duplicata duplicata* (Martin), BM R1366. Note that the cardinal septum extends across the corallite during early development (b-e) and that lonsdaleoid dissepiments of the parent are incorporated into the morphology of the daughter. 2 a-e, lateral increase in *L. floriformis* near *laticlavia* Smith, BM R36865. Note that the daughter corallite is nearly tangentially orientated to the parent. All figures  $\times 4$ .

Increase in *L. floriformis crassiconus* demonstrates particularly well the variability of development of minor septa. In some corallites, few if any minor septa are inserted until a late stage of development when major septa lengthen peripherally into the lonsdaleoid dissepiments during the late neanic stage (corallite D3 on Pl. 101, figs. 1 f-h); insertion is thus zaphrentid, followed by cyathaxonid. In other corallites, however, minor septa appear at an early stage of development after 10 to 15 major septa are inserted (Pl. 102, fig. 1 a). These minor septa disappear where lonsdaleoid dissepiments are formed by the withdrawal of major septa from the wall, and later reappear when major septa lengthen peripherally. The pattern of septal insertion during the hystero-ontogeny of these corallites is two cycles of zaphrentid insertion alternating with cyathaxonid insertion.

The pattern of septal insertion in *Lonsdaleia* appears to be dependent to some extent on individual variation and is thus of little taxonomic importance except that the earliest mode is always zaphrentid.

#### INCREASE IN *THYSANOPHYLLUM ORIENTALE*

*Thysanophyllum orientale*, chosen by Gregory (1917, p. 238) as type species of *Thysanophyllum*, occurs in the shale of Coral Zone 3 age at Aberlady Bay, Scotland. Hill (1940, p. 162) chose as lectotype specimen 1095 in the Thomson Collection, Kelvingrove Museum, Glasgow. This specimen is one of the two syntypes from Aberlady Bay on which Thomson (1880, p. 257) based the species. Only the two specimens were known from the type locality until recently, when two additional ones were collected. One of these, RSM 1964.25, which was collected by Dr. C. D. Waterston and Mr. I. F. Sime,





is dolomitized and examination of it could only be made on polished faces. This specimen differs from the lectotype in possessing slightly larger corallites, averaging 20 mm. in diameter, and more numerous septa, numbering about 30 of each order. Part of the early and late neanic stages of development were examined in two corallites and these resemble the characters of increase described below.

The other recently collected topotype, UQ F46767, which was collected by Mr. C. P. McDonagh, yielded good serial sections of parts of the hystero-ontogeny of six corallites (text-fig. 5). Corallites in this specimen (Pl. 102, figs. 5*a*, *b*) are slightly larger and have a few more septa than those in the lectotype. A prominent long axial septum, which Hill (1940, p. 162) regarded as the counter septum but in all likelihood is the cardinal septum, is present in a few adult or nearly adult corallites, whereas in the lectotype only some of the youthful corallites are known to possess this long septum. Finally, some corallites in UQ F46767 have a small imperisistently developed and isolated axial structure (text-fig. 5, 3*d-e*), a character unknown in the lectotype.

The other specimen studied by means of serial sections, RSM 1966.6.4, from Bathgate, West Lothian, resembles the corallum from this locality illustrated by Hill (1940, pl. 8, figs. 30-32). The hystero-ontogeny of seven corallites was studied in this specimen, but crushing has destroyed the initial stages of three of them and part of that in a fourth corallite.

Corallites arise by lateral increase in the peripheral part of the lonsdaleoid dissepimentarium of the parent. Increase normally starts with the appearance of five septa in the lonsdaleoid dissepimentarium of the parent (text-fig. 5, 1*b*, 3*a*). These septa correspond to the septa in the tabularium of the parent but are not united to them and they often do not terminate peripherally against lonsdaleoid dissepiments. The wall dividing daughter corallite from parent is formed near the peripheral ends of these septa, and during or following its formation these septa usually disappear.

*Hystero-brephic stage* (text-fig. 5, 1*d-e*). The cardinal septum is the first septum formed and the counter septum may appear shortly thereafter, or if the dividing wall is slow in being formed, it may not appear until after some metasepta have been inserted in the cardinal quadrants (text-fig. 5, 3*a*). Both axial septa are short but the cardinal septum is the longer of the two. In UQ F46767, this septum remains long throughout much of early development (text-fig. 5, 1*e-i*) but in RSM 1966.6.4 it soon shortens to equal the other septa in length. The axial plane is approximately radially orientated with respect to the axis of the parent. Alar septa may or may not be formed before metasepta appear, and are of the same length as metasepta during development. In UQ F46767 alar septa

---

TEXT-FIG. 5. Lateral increase in *Thysanophyllum orientale* Thomson, topotype, UQ F46767. Numbers below figures are cumulative distances of distal growth in mm. C = cardinal septum; K = counter septum; A = alar septum. 1*a-j*, typical hystero-ontogenetic sequence; *a-c*, initial stage of development, showing formation of five parent septa (*b*) and subsequent formation of the dividing wall (*c-d*); *d-e*, hystero-brephic stage; *f-j*, early neanic stage,  $\times 4$ . 2*a-e*, non-typical hystero-ontogenetic sequence in which the dividing wall is formed independently of septa and the parent corallite becomes incorporated into an adjacent adult corallite. Note that lonsdaleoid dissepiments are present at the beginning of development of the daughter corallite (*a-c*) and these subsequently become modified to interseptal dissepiments (*d-e*),  $\times 4$ . 3*a-e*, variation in the length of axial septa distally in an adult corallite and development of two daughter corallites,  $\times 2$ .

---

are positioned nearest the counter septum, but in RSM 1966.6.4 they are nearer the cardinal septum.

*Early neanic stage* (text-fig. 5, 1*f-j*). Insertion during this stage of development is not always confined to the fossulae and is quite variable, with groups of septa seeming to appear almost simultaneously. Minor septa, along with the first dissepiments, are formed in all quadrants at the same time as the major septa. Dissepiments during this stage of development are interseptal in nature, and about two or three rows are formed. Occasionally a few lonsdaleoid dissepiments are also present in parts of some corallites.

*Late neanic stage* (lower daughter corallite of text-fig. 5, 3*d*). The start of this stage is taken at the appearance of a lonsdaleoid dissepimentarium by the retreat of septa from the wall. The lonsdaleoid area of UQ F46767 is variably developed and some corallites may reach adult proportions and still lack a zone of these dissepiments around the corallite. Minor septa either disappear or become very short with the formation of the lonsdaleoid dissepiments.

A long cardinal septum and short counter septum are present in some adult corallites of UQ F46767 (figs. 3 *a-e*). Tracing these septa distally has shown that their length is quite variable, but shortening of the counter septum is more persistent than lengthening of the cardinal septum. The cardinal septum at its greatest length extends to the axis or slightly across it (fig. 3*c*), and when it is shortened, it may dissociate from its axial end leaving a thin axial structure (figs. 3 *d-e*). During further corallite growth, this axial structure slowly decreases in length and eventually disappears.

One unusual example of increase was observed in UQ F46767 (text-fig. 5, 2*a-e*), in which the daughter corallite arises in an aseptate parent and the dividing wall is formed without the usual association of parent septa. As this wall develops, the wall separating the parent corallite from an adjacent adult corallite is progressively shortened and the area formerly occupied by the parent becomes incorporated into this adjacent corallite.

#### COMPARISON OF CHARACTERS OF INCREASE

Significant differences in the hystero-ontogenies of *Lonsdaleia* and *Lithostrotion* (see Jull 1965) are as follows: (1) more corallites may develop simultaneously from the same parent in *Lonsdaleia* than is known in *Lithostrotion*; (2) the daughter and parent corallites of both cerioid and fasciculate species of *Lonsdaleia* are separated by a dividing wall, whereas in fasciculate forms of *Lithostrotion* a partition usually separates daughter from parent corallite; (3) *Lonsdaleia* has zaphrentid insertion in all quadrants during early development and a prominently long cardinal septum; (4) tangential orientation of daughter to parent occurs in some corallites of *Lonsdaleia* but is unknown in lateral increase in *Lithostrotion*; (5) the axial structure in *Lonsdaleia* is initiated by the axial end of the cardinal septum forming the median lamella, whereas in *Lithostrotion* the columella is usually formed by dilation at the axis of the united axial septa.

A comparison of the hystero-ontogeny of *Thysanophyllum orientale* with that in the above genera shows that this species is not closely allied to either genus. Some characters, however, suggest that it lies with the Lonsdaleiidae. The cardinal septum during early development resembles that in *Lonsdaleia* and the manner of formation of dissepiments

---

is fundamentally similar in both forms although they tend to appear later during development in *T. orientale*. Moreover, the axial structure in parts of some corallites of the topotype studied and of the specimen from Bathgate, West Lothian, which was figured by Hill (1940, pl. 8, fig. 31) is similar to that in diphymorphic corallites of *Lonsdaleia* (see Pl. 100, fig. 1). One character which might suggest affinities between *T. orientale* and the Lithostrotionidae is the mesa-shaped tabulae in the species, a character sometimes present in *Lithostrotion* but unknown in *Lonsdaleia*. Strikingly different from both *Lonsdaleia* and *Lithostrotion* is the irregular pattern of septal insertion and the appearance of minor septa in all quadrants at or near the start of insertion in *T. orientale*.

Unlike *Thysanophyllum minus* Thomson, which is most likely a diphymorph of *Lonsdaleia floriformis floriformis*, *T. orientale* is obviously not closely allied to *Lonsdaleia* but possibly shares a common ancestor with this genus. Smith (1916, p. 235) agreed with the suggestion of Carruthers (in Garwood 1913, p. 563) that the Lower Viséan  $C_1$  to  $C_2S_1$  *T. pseudovermiculare* McCoy is ancestral to *Lonsdaleia*. Perhaps it also gave rise to *T. orientale*. The horizon of the *T. pseudovermiculare* beds in the North-West Province is now considered to be wholly within the basal Viséan Upper Caninia ( $C_2S_1$ ) Zone (see Turner 1950, pp. 125, 128–9; 1959, p. 259).

In conclusion, *Thysanophyllum*, as presently interpreted, is undoubtedly a polyphyletic genus with some species related to the Lithostrotionidae. However, the family affinities of the genus must nevertheless lie with its type species, a probable lonsdaleoid, and species with different affinities can only be regarded as homeomorphs.


## REFERENCES

- DE GROOT, G. E. 1963. Rugose corals from the Carboniferous of Northern Palencia (Spain). *Leid. geol. Meded.* **29**, 1–123, 26 pl.
- DOBROLYUBOVA, T. A. 1958. Lower Carboniferous colonial tetracorals from the Russian Platform. *Trudy. paleont. Inst.* **70**, 1–224, 38 pl. [in Russian].
- EASTON, W. H. 1960. Permian corals from Nevada and California. *J. Paleont.* **34**, 570–83.
- FEDOROWSKI, J. 1965. Lower Permian Tetracoralla of Hornsund, Vestspitsbergen. *Studia geol. pol.* **17**, 1–173, 15 pl.
- FONTAINE, H. 1961. Les Madréporaires Paléozoïques du Viêt-Nam, du Laos et du Cambodge. *Archs géol. Viêt-Nam*, **5**, 276 pp., 35 pl.
- GARWOOD, E. J. 1913. The Lower Carboniferous succession in the north-west of England. *Q. Jl geol. Soc. Lond.* **68** (for 1912), 449–586, pl. 44–56.
- GREGORY, J. W. 1917. Thomson's genera of Scottish Carboniferous corals. *Trans. geol. Soc. Glasg.* **16**, 220–43.
- HILL, D. 1938–41. A monograph on the Carboniferous rugose corals of Scotland. *Palaeontogr. Soc. [Monogr.]*, 213 pp., 11 pl.
- 1956. Rugosa; In *Treatise on Invertebrate Paleontology* (Ed. R. C. MOORE). Part F, *Coelenterata*, pp. 233–324. Univ. Kansas Press and Geol. Soc. Am.
- JULL, R. K. 1965. Corallum increase in *Lithostrotion*. *Palaeontology*, **8**, 204–25.
- LECOMPTE, M. 1952. I. Madréporaires Paléozoïques; *Traité de Paléontologie* (Ed. J. PIVETEAU). Part 1, 419–539. Paris.
- MINATO, M. 1955. Japanese Carboniferous and Permian corals. *J. Fac. Sci. Hokkaido Univ.* (Ser. 4), **9**, 1–202, 43 pl.
- SMITH, S. 1916. The genus *Lonsdaleia* and *Dibunophyllum rugosum* (McCoy). *Q. Jl geol. Soc. Lond.* **71** (for 1915), 218–72, pl. 17–21.


- SOSHKINA, E. D. and DOBROLYUBOVA, T. A. 1962. Order Evenkiellida; In *Principles of Palaeontology* (U. A. Orlov, Ed.). *Spongia, Archaeocyatha, Coelenterata, Vermes* (Ed. B. S. Sokolov), 333-9. Acad. Sci. U.S.S.R. [in Russian].
- THOMSON, J. 1880. Contributions to our knowledge of rugose corals, from the Carboniferous Limestone of Scotland. *Proc. phil. Soc. Glasg.* **13**, 225-61, pl. 1-3.
- TURNER, J. S. 1950. Notes on the Carboniferous Limestone of Ravenstonedale, Westmorland. *Trans. Leeds geol. Ass.* **6**, 124-34.
- 1959. Tournaisian beds in Ravenstonedale and Dublin. *Geol. Mag.* **96**, 259-60.

R. K. JULL  
Department of Geology,  
University College of Townsville,  
Townsville, Queensland


Manuscript received 17 August 1966




1




2a  
0.00




2b  
0.60




2c  
1.40




2d  
1.64




2e  
1.88




2f  
2.14




2g  
2.44




2h  
2.58




2i  
2.74




2j  
2.94











2k  
3.26




2l  
3.58




2m







1a  
0.00



1b  
0.64




2



1c  
4.40



3



5a



5b



4a



4b