

WENLOCK STROPHEODONTIDAE (SILURIAN BRACHIOPODA) FROM THE WELSH BORDERLAND AND SOUTH WALES

by M. G. BASSETT

ABSTRACT. Eighteen species and subspecies of stropheodontid brachiopods are described from the Wenlock Series (Silurian) of the Welsh Borderland and South Wales, and are assigned to nine separate genera and subgenera. One new subgenus, *Amphistrophia* (*Pembrostrophia*), is recognized, together with the following new species and subspecies: *Strophonella* (*Strophonella*) *euglypha gentilis*, *Amphistrophia* (*Pembrostrophia*) *freshwaterensis*, *Mclearnites coralli*, and *Pholidostrophia* (*Mesopholidostrophia*) *deflecta*. Two separate stocks, represented by *Leptostrophia* (*Eostropheodonta*) and *Amphistrophia* (*Pembrostrophia*), are shown to retain dental plates into the Wenlock period.

ALTHOUGH the taxonomy and phylogenetic relationships of stropheodontid brachiopod genera have become fairly well established during the past thirty years, the recognition, interpretation, and nomenclature of British Wenlock species are still dependent largely on Davidson's Silurian Monograph (1866–71), in which all the strophomenid species were described in 1871. The purpose of the present paper is to systematically revise those stropheodontid species which occur in the Wenlock Series of the Welsh Borderland and South Wales in order to clarify their affinities with species from other areas and stratigraphical horizons. Material for this study has been collected mainly by the writer from the following areas: Wenlock Edge and Benthall Edge in Shropshire, Dudley and Walsall in the west English Midlands, the Ludlow Anticline, the Abberley and Malvern Hills, the inliers of May Hill, Woolhope, Usk, Tortworth, and Rumney (Cardiff), the Llandovery–Llandeilo district of Carmarthenshire, and sections at Freshwater East, Marloes Bay, and Winsle in Pembrokeshire. Supplementary collections have been studied in several museums in Britain, particularly the Davidson collection at the British Museum (Natural History) and the collections of the Geological Survey Museum, Institute of Geological Sciences.

Repositories. Throughout the text the repositories of type and figured specimens will be referred to by the following abbreviations: BM—British Museum (Natural History); GSM—Geological Survey Museum, Institute of Geological Sciences; NMW—National Museum of Wales, Cardiff; CMB—City Museum, Bristol; UB—University of Bristol Museum; SM—Sedgwick Museum, University of Cambridge; BU—Birmingham University (Geological Department Museum); NMI—National Museum of Ireland, Dublin; SMNH—Naturhistoriska Riksmuseet, Stockholm (Swedish Museum of Natural History); PMO—Paleontologisk Museum, Oslo.

In the Welsh Borderland the Wenlock Series comprises three formations, namely, in ascending order, the Woolhope Limestone, the Wenlock Shale, and the Wenlock Limestone. At Tortworth and throughout South Wales this tripartite division cannot

be recognized, the succession consisting instead of calcareous and sandy mudstones with thin limestone and sandstone horizons. The sequences throughout the area have recently been summarized elsewhere in tabular form (Bassett 1970, text-fig. 1) and their detailed correlation is currently being studied by the writer.

SYSTEMATIC DESCRIPTIONS

Order STROPHOMENIDA Öpik 1934

Superfamily STROPHOMENACEA King 1846

Family STROPHEODONTIDAE Caster 1939

The present supra-generic classification of the Stropheodontidae (Williams 1965, pp. H395–H403) is based mainly on the work of Caster (1939) and Williams (1953a). The latter author in 1953 established the use of subgenera to indicate the temporal relationships of evolving stropheodontid lineages and introduced the prefixes 'Eo', 'Meso', and 'Talaio' to denote early, mature, and late stages respectively. As more material is being examined in detail it is becoming evident that although the generally accepted picture of stropheodontid phylogeny still holds broadly true, the changes that take place in lineages of this type are not direct, and that early but conservative members of a stock may coexist for some time with their immediate descendants. For example Cocks (1967) showed that in the Upper Llandovery a number of forms with dental plates ('Eo' stage) may exist contemporaneously with direct descendants which lack dental plates ('Meso' stage). The same type of relationship can now be demonstrated with some of the Wenlock genera described below, and although Williams (1953a, p. 9 and text-fig. 4) considered that dental plates were lost in stropheodontids by the end of Llandovery times, at least two separate stocks can be shown to retain these structures well into the Wenlock period.

Subfamily STROPHEODONTINAE Caster 1939

Genus BRACHYPRION Shaler 1865

Type species. Strophomena leda Billings 1860, by original designation of Shaler 1865, p. 63.

Williams (1953a, p. 35; 1965, p. H395) regarded *Brachyprion* as a subgenus of *Strophodonta*, whereas both Cocks (1967, p. 256) and Havlíček (1967, p. 139) described it as an independent genus. The relatively gently concavo-convex lateral profile, and the configuration of the non-flabellate ventral muscle field in *Brachyprion* together indicate an affinity with the leptostrophiid stock, and the work of Cocks (1967, p. 258) suggests that it may have been derived from *Leptostrophia* s.s. in Middle or early Upper Llandovery times. In contrast, *Strophodonta* has a fairly strongly concavo-convex lateral profile and a large, well impressed, flabellate ventral muscle field, features which are together suggestive of a close relationship with the megastrophiiids. These postulated differences in the relationships of *Brachyprion* and *Strophodonta* are here considered sufficient to warrant their recognition as separate genera, but *Brachyprion* is retained as an early member of the Stropheodontinae, pending an investigation of its constituent species.

Brachyprion waltonii (Davidson)

Plate 54, figs. 1-5

- 1848 *Leptaena waltonii* Davidson, p. 317, pl. 3, fig. 6.
 1866 *Leptaena waltoni* Davidson; Davidson, pl. 8, fig. 19.
 1871 *Strophomena waltoni* (Davidson); Davidson, p. 310, pl. 42, fig. 11.
 1956 *Strophomena waltoni* (Davidson); Curtis, p. 150.
 1967 *Brachyprion waltoni* (Davidson); Cocks, p. 256.

Description. Exterior. Gently to moderately concavo-convex with subquadrate outline. Hinge-line straight, cardinal angles obtusely rounded. Point of maximum width commonly at about mid-length of shell. Lateral margins gently curved, anterior margin evenly rounded or sub-parallel to hinge. Anterior commissure rectimarginate.

Ventral interarea plane, apsacine, 2 to 3 times as long as dorsal interarea. Presence or absence of pseudodeltidium not apparent on material available. Delthyrial angle about 80°. Dorsal interarea short, anacline. Notothyrium sealed by massive, convex chilidium which extends up into delthyrium.

Radial ornament equally to subequally parvicostellate with an average of 4 costellae per mm over whole surface of shell. Costellae are low, rounded, and radiate from umbo with little curvature. Very fine, closely spaced growth fila are visible on well-preserved specimens.

Interior of pedicle valve. Ventral process low, produced anteriorly as slender myophragm which longitudinally bisects muscle field. Extent of denticulation of hinge not observed. Muscle field broadly triangular in outline, moderately well impressed, confined to posterior half of valve, bounded laterally by pair of low ridges which are antero-laterally divergent at about 80-90° to one another. Fairly coarse pseudopunctae cover interior surface of valve, apart from muscle field.

Interior of brachial valve. Unknown.

Dimensions of figured specimens in mm

		Length	Maximum width	Hinge width
Complete shell	Holotype, BM B5989	18.9	22.3	18.0
Complete shell	UB 12325-1	19.4	21.0	18.3
Partially exfoliated exterior of pedicle valve	UB 12325-2	19.5	22.6	19.0 est.
Partially exfoliated exterior of pedicle valve	GSM DEW9625	16.4	19.6	16.1
Impression of exfoliated pedicle valve	GSM DEW9627	—	—	—

Holotype (by monotypy). BM B5989, from the 'Wenlock Shale, Falfield', Tortworth inlier, Glos. This is the specimen that was originally figured by Davidson (1848, pl. 3, fig. 6) and later refigured by him (1866, pl. 8, fig. 19; 1871, pl. 42, fig. 11). It is refigured on Plate 54, figs. 1a-c. The Wenlock Shale horizon at Falfield probably refers to the *Pyenactis* band which immediately overlies the limestone band at the base of the Wenlock Series (Curtis 1955, p. 6; 1956, p. 152).

The holotype of *B. waltonii* is rather unusual in that it has attached to it the 3 syntype specimens of the inarticulate brachiopod '*Crania*' *siluriana* Davidson 1866, p. 82, plate 8, figs. 19, 20. These 3 specimens bear a separate registration number (BM 15034-6).

Remarks. In synonymy lists of this species Davidson twice (1848, p. 317; 1871, p. 310) referred to *Leptaena waltonii* Davidson as being published in the *London Geological*

Journal for 1847, plate 26, fig. 3. However, plate 26 was never published in that issue of the journal and the first description and figure of *L. waltonii* were not published by Davidson until 1848. The original spelling of the specific name was *waltonii*, although in later papers Davidson and other authors used only a single *i* (see synonymy). Recommendation 31A of the International Code of Zoological Nomenclature (1964) states that there is no mandatory spelling for specific names based on modern personal names, although the single *i* is preferred. It is now usual to keep the double *i* in those names originally spelt in that way but to use the single *i* for new names. The original spelling of *waltonii* should therefore be maintained.

Distribution. *B. waltonii* is known only from the Tortworth inlier, Gloucestershire. As noted above the holotype is probably from the *Pycnactis* band which immediately overlies the limestone band at the base of the Wenlock Series; all other specimens examined by the writer are from the basal limestone of Brinkmarsh Quarry, 400 yd NNW of Brinkmarsh Farm, Whitfield (Grid Ref. ST 6735.9130).

Brachyprion sp.

Plate 54, figs. 6–8

Comparison. *Brachyprion* sp. is distinguished from *B. waltonii* chiefly by its unequally parvicostellate ornament and relatively wider hinge-line, which commonly forms the point of maximum width of the shell. In *B. waltonii* the ornament is equally to subequally parvicostellate and the point of maximum width is near to the mid-length.

EXPLANATION OF PLATE 54

Figs. 1–5. *Brachyprion waltonii* (Davidson).

1 *a–c.* Dorsal, ventral, and lateral views of complete shell, holotype, BM B5989, $\times 1.3$, 'Wenlock Shale' (probably *Pycnactis* band), Falfield, Tortworth inlier, Glos.

2*a, b.* Dorsal and ventral views of complete shell, UB 12325–1, $\times 1.3$, basal Wenlock (*Pycnactis* band or basal limestone band), Brinkmarsh Quarry, Whitfield, Tortworth inlier, Glos., ST/6735.9130.

3, 4. Partially exfoliated exterior of pedicle valve; 3, UB 12325–2; 4, GSM DEW9625, $\times 1.3$, basal limestone of Wenlock Series, locality as for fig. 2.

5. Impression of exfoliated pedicle valve, GSM DEW9627, $\times 1.3$, basal limestone of Wenlock Series, locality as for fig. 2.

Figs. 6–8. *Brachyprion* sp. Wenlock Shale, railway cutting, Daw End, Walsall, Staffs., SK/0360.0030.

6*a–c.* Dorsal, ventral, and lateral views of broken shell, NMW 70.3G.1, $\times 1.3$.

7*a, b.* Dorsal and ventral views of broken shell, NMW 70.3G.2, $\times 1.3$.

8. Artificially produced (calcined) internal mould of brachial valve, NMW 70.3G.3*a*, $\times 1.3$, from approx. 200 ft below Wenlock Limestone.


Figs. 9–13. *Megastrophia* (*Protomegastrophia*) *semiglobosa* (Davidson).

9*a, b.* Ventral and lateral views of pedicle valve, lectotype, BM B13642, $\times 1$, Wenlock Shale or Limestone, Rushall Canal, near Walsall, Staffs.

10, 13. Interiors of brachial valves; 10, BM B45075; 13, BM 746 $\times 1$, Wenlock Limestone, Dudley, Worcs.

11. Interior of pedicle valve, BM BB6419, $\times 1$, Wenlock Limestone, Dudley, Worcs.

12. Internal mould of pedicle valve, GSM 12758, $\times 1$, Wenlock Shale, Usk inlier, Mon.



BASSETT, Wenlock Stropheodontidae

The following description of *Brachyprion* sp. is based on a collection of 18 incomplete articulated specimens, a number of which have been roasted in a Bunsen flame and the calcined shell material scraped away to produce artificial internal moulds. The detail that can be observed in preparations of this type is rather limited and while an assignment to *Brachyprion* is possible, based on the convexity of the valves, the musculature, and the nature of the ornament, specific determination is not thought advisable until more material becomes available and related species can be compared in detail.

Description. Exterior. Shell thin, gently concavo-convex, body cavity narrow. Subquadrate to semicircular in outline with maximum width commonly at straight hinge-line, which may be produced as short, rounded ears. Lateral and anterior margins evenly curved; anterior commissure rectimarginate.

Ventral interarea plane to very gently concave, apsacline, 4–5 times as long as dorsal interarea, which is plane, anacline. Median third of both interareas bears fine ridges caused by impress of internal denticulation. Delthyrial angle about 80°. Pseudodeltidium small, gently convex; chilidium massive, convex, completely sealing notothyrium and extending up into delthyrium.

Ornament unequally parvicostellate with up to 4 fine costellae occupying interspaces between larger, rounded costellae. There are 4–5 costellae per mm at the 5-mm growth stage of pedicle valve. Costellae increase in strength anteriorly, are straight along mid-line of shell and curve gently on flanks. Whole surface is crossed by extremely fine, concentric growth fila.

Interior of pedicle valve. Pseudodeltidium supported by low, short ventral process, gently concave along its mid-length and extending anteriorly into shallow, narrow, rounded groove bisecting muscle field. Process pits small, shallow. Muscle field broadly triangular in outline, occupying about one-third of valve length and up to one-third width. Muscle field unbounded anteriorly but may be bounded laterally by low ridges. Adductor scars narrow, elongate, situated adjacent to median groove and confined to posterior half of muscle field. Diductor scars large, sub-triangular, enclosing adductor scars laterally and anteriorly.

Interior of brachial valve. Cardinal process lobes narrow, platelike, initially divergent at about 160° to one another but curving anteriorly to be finally divergent at about 90°. Socket ridges well developed, antero-laterally divergent at about 100° to one another. Notothyrial platform low, hollowed out immediately below cardinal process lobes and produced anteriorly as low, convex ridge which longitudinally bisects muscle field.

Muscle field subquadrate, well impressed, occupying about one-third of valve length and one-quarter width; laterally bounded by broad, high, slightly curved ridges, but remains unbounded anteriorly. Pair of lanceolate scars lies alongside median ridge, enclosed by larger, more deeply impressed pair of scars with sub-oval outline.

Internal surface of both valves covered with fine, radially aligned pseudopunctae, and shows impress of external ornament.

Distribution. *Brachyprion* sp. is known only from the Wenlock Shale of the railway cutting at Daw End, Walsall, Staffordshire (Grid Ref. SK 0360.0030).

Genus MEGASTROPHIA Caster 1939
 Subgenus *Megastrophia* (*Protomegastrophia*) Caster 1939

Type species. *Leptaena profunda* Hall 1852, by original designation of Caster 1939, p. 28.

Megastrophia (*Protomegastrophia*) *semiglobosa* (Davidson)

Plate 54, figs. 9–13; Plate 55, figs. 1–3

- 1847 *Leptaena imbrex* (Pander); Davidson, p. 55, pl. 12, figs. 25–8; non Pander 1830.
 1848 *Leptaena imbrex* (Pander); Davidson, p. 318, pl. 3, fig. 8; non Pander 1830.
 1854 *Strophomena imbrex* (Pander) Salter in Murchison, table on p. 488, woodcut 41, p. 224, figs. 6, 7; non Pander 1830.
 1871 *Strophomena imbrex* (Pander) var. *semiglobosa* Davidson, p. 286, pars, pl. 41, figs. 1–4, non figs. 5, 6.
 1883 *Strophomena imbrex* var. *semiglobosa* Davidson; Davidson, p. 195.
 1953a *Megastrophia semiglobosa* (Davidson); Williams, p. 21.

Comparison. Cocks (1967, p. 260) erected a new subgenus, *Eomegastrophia*, to include megastrophids which possess dental plates. The type of this subgenus is *M. (E.) ethica* Cocks from the Upper Llandovery of Shropshire and is probably directly ancestral to the Wenlock species described here. While the two species are closely related in details of musculature, ornament, and convexity, *M. (P.) semiglobosa* may easily be distinguished by the absence of dental plates. The Scottish Ordovician specimens figured by Davidson (1871, pl. 41, figs. 5, 6) are not stropheodontids; they belong to *Dactylogonia? semiglobosina* (Davidson) (see Davidson 1883, pp. 195–6; Williams 1962, pp. 201–2).


EXPLANATION OF PLATE 55

Figs. 1–3. *Megastrophia* (*Protomegastrophia*) *semiglobosa* (Davidson).

- 1a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.4, $\times 1$, Wenlock Shale, exposures below tumuli on Craig-y-Garced, 320 yd NW of Hen-rhiw Farm, Usk inlier, Mon., SO/3570.0275.
 2. Internal mould of pedicle valve, GSM 12756, $\times 1$, Wenlock Shale, Usk inlier, Mon.
 3. Internal mould of brachial valve, GSM HC991, $\times 1$, late Wenlock mudstone, old overflow trench near Middleton Hall, Carms., SN/5265.1879.

Figs. 4–11. *Strophonella* (*Strophonella*) *euglypha* (Dalman).

- 4a–d. Dorsal, ventral, lateral and posterior views of complete shell, BU 405 (ex Ketley collection), $\times 1$, Wenlock Limestone, Dudley, Worcs.
 5. Exterior of brachial valve, SMNH Br2342, lectotype, $\times 1$, Mulde Marl, Djupvik, Gotland.
 6. Interior of pedicle valve, BU 406 (ex Ketley collection), $\times 1$, Wenlock Limestone, Dudley, Worcs.
 7a, b. Dorsal and posterior views of interior of brachial valve, BM BB32805 (ex B5675), $\times 1$, Wenlock Limestone, Dudley, Worcs.
 8. Interior of brachial valve, BU 407 (ex Ketley collection), $\times 1$, Wenlock Limestone, Dudley, Worcs.
 9. Internal mould of brachial valve, GSM 13369, $\times 1$, Aymestry Limestone (Ludlow Series), Dormington Wood, probably Woolhope inlier, Herefordshire.
 10. Internal mould of pedicle valve, NMW 70.3G.5a, $\times 1$, siltstone overlying Wenlock Limestone, old quarry at Cwm, Usk inlier, Mon., SO/3331.0160.
 11 a–c. Dorsal, ventral, and lateral views of complete shell, SMNH Br33324, $\times 1$, Gotland, probably Mulde Marl, Djupvik.



BASSETT, Wenlock Stropheodontidae

Description. Exterior. Fairly large, thick shells, moderately to strongly concavo-convex, sometimes flattened slightly towards cardinal extremities. Outline subcircular with maximum width at any point between hinge and mid-length of shell. Hinge-line straight, commonly produced as short, rounded ears. Lateral margins commonly evenly curved, but may show slight constriction immediately anterior to hinge. Commissures smooth, anterior commissure rectimarginate.

Ventral interarea plane, apsacline, up to 3 times as long as dorsal interarea. Delthyrial angle varies from about 80–100°. Pseudodeltidium small, apical, gently convex. Dorsal interarea plane, anacline. Notothyrium closed by massive, convex chilidium which extends up into delthyrium.

Ornament unequally parvicostellate. Up to 8 fine, threadlike costellae occupy broad interspaces between stronger, rounded costellae. Costellae fairly straight, about 4 or 5 per mm at 5-mm growth stage of pedicle valve.

Interior of pedicle valve. Ventral process low, gently concave along its mid-length, produced anteriorly as slender myophragm. Anteriorly, myophragm decreases in height and width and in some specimens passes into shallow groove. Process pits shallow. Hinge-line denticulate for up to one-third of its length, denticles being borne on pair of low denticular plates.

Muscle field broadly triangular in outline, becoming flabellate anteriorly. Bounding ridges low or absent. Adductor scars small, narrow, lanceolate, situated in centre of muscle field and separated by myophragm; bounded laterally and anteriorly by large, sub-oval diductor scars. Area immediately lateral to muscle field may be coarsely pustulose, and whole of internal surface of valve bears fairly coarse pseudopunctae.

Interior of brachial valve. Cardinalia are supported on broad, low notothyrial platform which is deeply hollowed out below cardinal process lobes; these lobes consist of a pair of stout, upright plates with attachment faces directed postero-laterally, and are separated by rounded groove bearing low, triangular ridge. Socket plates weak, flat, widely divergent antero-laterally, with crenulations on their posterior faces.

Muscle field elongate-oval in outline, consisting of 2 oval adductor scars separated by prominent myophragm which narrows anteriorly. Muscle field occupies up to one-third of valve width and up to one-half length.

Median third of hinge bears denticles complementary to those of pedicle valve.

Dimensions of figured specimens in mm

		Length	Width
Pedicle valve	Lectotype, BM B13642	51.5	—
Interior of pedicle valve	BM BB6419	26.0	32.8
Internal mould of pedicle valve	GSM 12756	24.8	33.6 est.
Internal mould of pedicle valve	GSM 12758	34.0 est.	—
Interior of brachial valve	BM B746	42.6	45.6
Interior of brachial valve	BM B45075	38.7	42.9
Internal mould of pedicle valve	NMW 70.3G.4	33.4	45.0
Internal mould of brachial valve	GSM HC991	38.5	—

Lectotype (here selected): BM B13642 (figured Davidson 1871, pl. 41, fig. 2). This is one of 4 specimens in the Davidson collection which are mounted on a card bearing Davidson's handwriting and identifying the specimens as types. The other 3 specimens cannot be identified with certainty with Davidson's

figures, which were often composite, but his plate 41, figs. 1 *a-b*, are possibly based on the same collection. The card in Davidson's handwriting states that the types are from the Wenlock Limestone of Walsall and Dudley. According to the plate legend, however, the lectotype is from the Wenlock Shale of Rushall Canal, near Walsall; its exact horizon is therefore in some doubt. The lectotype is refigured in Plate 54, figs. 9*a, b*.

Distribution. This species does not occur very commonly in present Wenlock outcrops but numerous specimens have been examined in museum collections from the Wenlock Shale and Limestone of Walsall and Dudley. It has been collected by the writer from the Wenlock Shale of the Usk inlier, the late Wenlock mudstones of the Rumney (Cardiff) inlier and the late Wenlock mudstones of the Llandeilo district, Carmarthenshire. It is not known with certainty from early Wenlock horizons and has not been found in Pembrokeshire.

Genus STROPHONELLA Hall 1879

Subgenus *Strophonella* (*Eostrophonella*) Williams 1950

Type species. *Amphistrophia davidsoni* Høltedahl 1916, by original designation of Williams 1950, p. 281.

Discussion. No Wenlock species of *Strophonella* (*Eostrophonella*) are known, the subgenus being recorded only from the Llandovery Series (e.g. see Williams 1951, p. 127). It is discussed here, however, since during the course of this work it became necessary to clarify the affinities of the poorly known species '*Strophomena*' *simulans* McCoy 1851; examination of the type specimens of *simulans* reveals that the original description was based on poorly preserved material containing, among others, a specimen of *Strophonella* (*Eostrophonella*) of probable Lower Llandovery age, and a specimen of *Strophonella* (*Strophonella*) from the Wenlock Series. The Wenlock specimen is redescribed in this paper as *S.* (*Strophonella*) *euglypha gentilis* subsp. nov. (see p. 313) and the specific name *simulans* is retained for the specimen of *S.* (*Eostrophonella*). The relationships of these specimens are discussed in full on p. 314 following the description of *S.* (*S.*) *euglypha gentilis*.

Subgenus *Strophonella* (*Strophonella*) Hall 1879

Type species. *Strophomena semifasciata* Hall 1863, by subsequent designation of Hall and Clarke 1892, p. 291.

Strophonella (*Strophonella*) *euglypha* (Dalman)

Plate 55, figs. 4-11; Plate 56, figs. 1, 2

- 1828 *Leptaena euglypha* Dalman, p. 108, pl. 1, fig. 3*a, c*, non fig. 3*b*.
- 1828 *Leptaena euglypha* Dalman; Hisinger, pl. 4, figs. 7*a, b*.
- 1839 *Leptaena euglypha* Dalman; J. de C. Sowerby in Murchison, p. 622, pl. 12, fig. 1.
- 1847 *Leptaena euglypha* Dalman; Davidson, p. 56, pl. 12, figs. 1-4.
- 1848 *Leptaena euglypha* Dalman; Davidson, p. 316, pl. 3, fig. 4.
- 1852 *Leptaena* (*Strophomena*) *euglypha* (Dalman); McCoy, p. 243.
- 1861 *Strophomena euglypha* (Dalman); Lindström, p. 372.
- 1871. *Strophomena euglypha* (Hisinger); Davidson, p. 288, *pars* pl. 40, figs. 1-3, non figs. 4, 5.
- ? 1916 *Strophonella euglypha* (Hisinger); Høltedahl, p. 66, pl. 6, figs. 5, 6.

- 1953a *Strophonella euglypha* (Hisinger); Williams, p. 21.
1963 *Strophonella euglypha* (Hisinger); Holland *et al.*, pl. 3, figs. 8, 9.
? 1967 *Strophonella euglypha* (Hisinger); Havlíček, p. 180, pl. 37, figs. 6–8; pl. 50, figs. 5, 8, 9.

Comparison. Williams (1951, p. 128) showed that two of the specimens figured by Davidson in 1871 (pl. 40, figs. 4, 5) as '*Strophomena euglypha*' belong to *Strophonella* (*Eostrophonella*) *davidsoni* (Holtedahl). This Llandovery species differs from *S. (S.) euglypha* in having short dental plates in the interior of the pedicle valve and in having a less prominent, more rounded ventral muscle field.

Description. Exterior. Pedicle valve initially gently convex; brachial valve initially plane to weakly concave. At length of about 13–20 mm valves become resupinate and develop strong convex-concave profile. Outline semicircular to subtriangular, varying from as wide as long to about seven-tenths as long as wide. Maximum width at or slightly posterior to straight hinge-line, which in well-preserved shells is produced as short, rounded ears. Margins of shell commonly evenly curved but anterior margin may be extended anteriorly as tongue-like trail to give an overall subtriangular outline. Commissures smooth, anterior commissure rectimarginate.

Ventral interarea plane, apsacline, 2–3 times as long as dorsal interarea. Delthyrial angle about 40°. Delthyrium almost completely closed by well-developed, convex pseudodeltidium. Dorsal interarea linear, plane, anacline. Chilidium strong, convex, completely sealing notothyrium. Both interareas show traces of internal denticulation.

Ornament unequally parvicostellate with up to 6 (commonly 2 or 3) very fine capillae occupying wide interspaces between major rounded costellae. At 10-mm growth stage there are 4–5 costellae per mm. Very fine, closely spaced, concentric growth fila may be seen on well-preserved shells. A number of specimens exhibit fine crenulations of shell along posterior margin, inclined obliquely to hinge (e.g. see Pl. 55, figs. 4a, b).

Interior of pedicle valve. Pseudodeltidium supported by stout, hastate, ventral process, hollowed out along its mid-length, and produced anteriorly as strong, rounded myophragm. Deep process pits laterally bound ventral process. Dental plates absent. Median third of hinge bears pair of well-developed denticular plates.

Muscle field strongly impressed, subquadrate, flabellate, occupying about one-third of width of valve and up to one-half length. Strong muscle bounding ridges arise immediately anterior to hinge, diverging initially at about 90° to one another, then becoming parallel and finally turning medially and posteriorly to intersect myophragm slightly posterior to its anterior extremity. Adductor scars narrow, lanceolate, situated adjacent to myophragm and surrounded by larger, oval diductor scars.

Extra-muscular area shows traces of external ornament and may be coarsely pustulose.

Interior of brachial valve. Cardinalia raised on broad notothyrial platform, produced anteriorly as convex median ridge. Cardinal process lobes stout, pear-shaped, divergent, grooved along their mid-length, with attachment faces directed postero-ventrally. Sockets shallow, socket plates short, strong, antero-laterally divergent at about 110° to one another.

Muscle field moderately well impressed, truncated oval in outline, occupying about one-quarter of valve width and commonly about two-thirds as long as wide. Low,

broad, slightly curved ridges may bound muscle field laterally but it is unbounded anteriorly and bisected longitudinally by median ridge which tapers anteriorly.

Internal surface may be coarsely tuberculate, especially immediately lateral to muscle field.

Dimensions of figured specimens in mm

		<i>Length</i>	<i>Width</i>
Exterior of brachial valve	Lectotype, SMNH Br2342	28.5	45.0 est.
Complete shell	SMNH Br33324	35.0	—
Interior of pedicle valve	SMNH Br33301	29.1	—
Complete shell	BU 405 (ex Ketley collection)	35.6	40.5
Interior of pedicle valve	BU 406 (ex Ketley collection)	36.4	38.1
Interior of pedicle valve	BU Holcroft Collection no. 297	39.6	40.5
Interior of brachial valve	BU 407 (ex Ketley collection)	—	34.9
Interior of brachial valve	BM BB32805 (ex B5675)	—	—
Internal mould of brachial valve	GSM 13369	35.8	32.5 est.
Internal mould of pedicle valve	NMW 70.3G.5a	—	41.3

Type and figured specimens. Lectotype (here selected): the specimen figured by Dalman 1828, plate 1, fig. 3a, now housed in the Department of Palaeozoology, Swedish Museum of Natural History, Stockholm (Br2342, Hisinger sample 472). This specimen, from the Mulde Marl of Djupvik, Gotland is refigured on Plate 55, fig. 8. The Mulde Marl is regarded as being of late Wenlock age (Hede 1960, p. 49). The specimen figured by Dalman 1828, plate 1, fig. 3b (SMNH Br2336, Hisinger sample 473) does not belong to *S. euglypha* but to the genus *Megastrophia*. Examination of the lectotype and possible topotype material of *S. euglypha* leaves no doubt that the British and Swedish shells are conspecific.

Remarks. The authorship of this well-known species has been the subject of some confusion in the past (see synonymy), being referred by various authors to either Dalman

EXPLANATION OF PLATE 56

Figs. 1, 2. *Strophonella (Strophonella) euglypha* (Dalman).

1. Interior of pedicle valve, BU Holcroft collection no. 297, $\times 1$, 'Upper Silurian' (probably Wenlock Limestone), Wren's Nest, Dudley, Worcs.

2. Interior of pedicle valve, SMNH Br33301, $\times 1$, Gotland, probably Mulde Marl, Djupvik.

Figs. 3–6. *Strophonella (Strophonella) euglypha gentilis* subsp. nov.

3 a–c. Internal mould of pedicle valve, latex cast, and counterpart external mould, holotype, NMW 70.3G.6a, b, $\times 1$, late Wenlock mudstone, forestry track section, W side of Sawdde gorge, Carms., SN/7200.2525.

4a, b. Internal mould of brachial valve and latex cast, NMW 70.3G.24a, $\times 1$, locality and horizon as for fig. 3.

5a, b. Internal mould of pedicle valve and counterpart external mould, SM A10272a, b, $\times 1$, Wenlock Series, Middleton Park, Carms.

6. Internal mould of pedicle valve, GSM DEX2853, $\times 1$, late Wenlock mudstone, old overflow trench near Middleton Hall, Carms., SN/5265.1879.

Figs. 7, 8. *Leptostrophia (Eostropheodonta)* sp.

7a, b. Internal mould of pedicle valve and counterpart external mould, NMW 70.3G.11a, b, $\times 1.3$, Wenlock Shale, exposures below tumuli on Craig-y-Garced, 320 yd NW of Hen-rhiw Farm, Usk inlier, Mon., SO/3570.0275.

8. External mould of pedicle valve, NMW 70.3G.12a, $\times 1.3$, locality and horizon as for fig. 7.


Figs. 9–10. *Leptostrophia (Leptostrophia) filosa* (J. de C. Sowerby).

9a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.9a, $\times 1.3$, siltstone overlying Wenlock Limestone, old quarry at Cwm, Usk inlier, Mon., SO/3331.0160.

10. Exterior of pedicle valve, GSM 13307, $\times 1.3$, Wenlock Limestone, Usk inlier, Mon.; figured Davidson 1871, pl. 44, fig. 16.



1



3a



3c




2




3b



4a



5a




5b




4b




6




7a




7b



8



9a



9b



10

BASSETT, Wenlock Stropheodontidae

or Hisinger and even to Sowerby by Davidson (1871, pl. 40) in his plate legend. Dr. V. Jaanusson has kindly supplied the writer with information to help to clarify the matter. Davidson's reference (1871, p. 288) to *Leptaena euglypha* Hisinger 1819 is certainly a mistake since the species is neither mentioned nor figured in Hisinger's paper of that year. The species name *Leptaena euglypha* was first published in the same year (1828) by both Dalman and Hisinger, though it is not known which publication has priority. However, it seems best to refer the species to Dalman since Hisinger does so and since Dalman does not mention Hisinger's authorship. The problem now is whether the author should be regarded as Dalman 1828 or Dalman *in* Hisinger 1828. This depends on which of the publications has priority and there may be a question of priority of days. Fortunately this is a purely academic matter because the description of the species in both papers is based on the same material and thus selection of the lectotype from the specimens figured by Dalman satisfies the nomenclatural requirement of both papers.

Distribution. *S. (S.) euglypha* is widespread in the Wenlock of the Welsh Borderland, occurring most commonly in the Wenlock Limestone and less commonly in the upper calcareous horizons of the Wenlock Shale. It is unknown to the writer from low in the Wenlock Shale but a single specimen is recorded from the Woolhope Limestone in the Geological Survey collections (GSM 12679). The species is also found in late Wenlock mudstones and sandstones at Tortworth and Rumney (Cardiff). In the Llandovery-Llandeilo district, Carmarthenshire, *S. (S.) euglypha* s.s. appears to be replaced by *S. (S.) euglypha gentilis* (see below); neither subspecies is known from Pembrokeshire.

Lawson (1960, p. 117) recorded *S. euglypha* as rare or absent in Britain after Bringerwoodian times, while in Gotland it was recorded by Hede (1960) from localities in the Mulde Marl (late Wenlock) and from the Hemse group (early Ludlow). Holteedahl (1916, p. 66) and Havilíček (1967, p. 180) recorded *S. euglypha* from Norway and Czechoslovakia respectively, though these identifications have not been checked. The full recorded range of the species is from early Wenlock to Ludlow (Bringerwoodian).

Strophonella (Strophonella) euglypha (Dalman) *gentilis* subsp. nov.

Plate 56, figs. 3-6

1851 *Strophomena simulans* McCoy, p. 403, *pars*.

1852 *Strophomena simulans* McCoy; McCoy *in* Sedgwick and McCoy, p. 246 *pars*, pl. 1H, fig. 35, *non* figs. 33, 34.

1871 *Strophomena simulans* McCoy; Davidson, p. 297, *pars*, pl. 42, fig. 10, *non* figs. 9, 9 a-c.

Discussion. Specimens of *Strophonella* that have been collected from the late Wenlock mudstones and siltstones of the Llandeilo district, Carmarthenshire, are closely similar in outline and ornament to *S. (S.) euglypha* from the more calcareous horizons of the Welsh Borderland. The South Wales specimens may, however, be distinguished by having a more rounded ventral muscle field, a less-strongly impressed dorsal muscle field, and by being relatively broader and more gently resupinate than is typical of *S. (S.) euglypha* s.s. These differences are probably due largely to the more clastic environment in the South Wales area, but the specimens are sufficiently distinct to warrant their recognition as a new subspecies within *S. (S.) euglypha*.

Dimensions of figured specimens in mm

		Length	Width
Internal mould of pedicle valve and counterpart external mould	Holotype, NMW 70.3G.6 a, b	38.3	60.0 est.
Internal mould of brachial valve	NMW 70.3G.24a	—	—
Internal mould of pedicle valve and counterpart external mould	SM A10272 a, b	—	—
Internal mould of pedicle valve	GSM DEX2853	—	60.8

Type and figured specimens. Holotype, NMW 70.3G.6a, b, the internal mould of a pedicle valve and its counterpart external mould, figured on Plate 56, figs. 3a–c. The other specimens whose dimensions are given above are paratypes. One of these (SM A10272a, b) was included by McCoy and Davidson (see synonymy) in '*Strophomena simulans* McCoy, but as the lectotype of *simulans* is a specimen of *Strophonella* (*Eostrophonella*), of probable Lower Llandovery age (see discussion below), the Wenlock specimen of *Strophonella* (*Strophonella*) is re-assigned here to the new subspecies *gentilis*.

Distribution. *S. (S.) euglypha gentilis* occurs fairly commonly in the late Wenlock mudstones of the forestry track section along the west bank of the Sawdde gorge, Carmarthenshire (Grid Ref. SN/7200.2525) where it is associated with the graptolite *Monograptus flemingii flemingii* Salter. It has also been found in sandy siltstones of Wenlock age in an old overflow trench near Middleton Hall, Carmarthenshire (Grid. Ref. SO/5265.1879). Large strophonellids collected from comparable horizons throughout the Llandeilo district are generally broken, but almost certainly belong to this subspecies.

Discussion. As discussed both immediately above and earlier (p. 310), McCoy's original description (1851, p. 403) of *Strophomena simulans* was based on a number of specimens now known to belong to different species. The same specimens were later figured by McCoy (1852, pl. 1H, figs. 33–35) and by Davidson (1871, pl. 42, figs. 9, 10). All the material is now in the Sedgwick Museum, Cambridge, and in order to clarify their relationships, each specimen is discussed separately.

1. SM A42416a, b. Internal mould of pedicle valve and counterpart external mould. Here selected as lectotype of *Strophomena simulans* and identified as a species of *Strophonella* (*Eostrophonella*). A42416a was figured by McCoy 1852, plate 1H, fig. 34 and in the reconstruction of fig. 34a. The specimens are from Cefn Coch, Glyn Ceiriog, Denbighshire ('Upper Bala' of McCoy); the locality suggests that they were from the Fron Frys Slates, of probable Lower Llandovery age (Groom and Lake 1908, p. 553, pl. 53; Curtis in Whittard 1961, p. 92).

2. SM A42417. Internal mould of pedicle valve of an indeterminate resupinate strophomenid, possibly a species of *Amphistrophia*. Used in the reconstruction of McCoy 1852, plate 1H, figs. 33, 34a. This specimen is from Blain y Cwm, west of Nantyre, Glyn Ceiriog, Denbighshire ('Upper Bala' schists of McCoy); the exact horizon is unknown.

3. SM A45163. External mould of brachial valve of a strophomenid, possibly *Megastrophia* (*Protomegastrophia*) *semiglobosa* (Davidson). Probably used to complete the restoration of McCoy 1852, plate 1H, figs. 33, 34a. The specimen is from McCoy's 'Upper Bala', Golden Grove, Llandeilo, Carmarthenshire; the locality is suggestive of a Wenlock age, a conclusion that is supported by the lithology and by the presence in the same slab of an eospiriferid, probably *Cyrtia exprorecta* (Wahlenberg).

4. SM A10272a, b. Internal mould of pedicle valve and counterpart external mould. This is the specimen identified and figured (Pl. 56, figs. 5a, b) as *S. (S.) euglypha gentilis*; it was figured by McCoy 1852, plate 1H, fig. 35. It is the specimen referred to by McCoy (1851, p. 404; 1852, p. 246) as 'one doubtful specimen in the sandy schists of the Malverns, Worcestershire'. The locality is almost certainly a mistake, since both Salter (1873, p. 147) and Woods (1891, p. 64) indicated that it was collected

from the Wenlock of Middleton Park, Carmarthenshire. The lithology of the specimen compares closely with that in material collected by the writer from localities in Middleton Park, while neither the subspecies nor similar lithological types have been identified in numerous collections from the Malverns.

Subfamily LEPTOSTROPHIINAE Caster 1939

Genus LEPTOSTROPHIA Hall and Clarke 1892

Subgenus *Leptostrophia* (*Leptostrophia*) Hall and Clarke 1892

Type species. *Stropheodonta magnifica* Hall 1857, by original designation of Hall and Clarke 1892, p. 288.

Leptostrophia (*Leptostrophia*) *filosa* (J. de C. Sowerby)

Plate 56, figs. 9, 10; Plate 57, figs. 1-6

- 1839 *Orthis filosa* J. de C. Sowerby in Murchison, p. 630, pl. 13, fig. 12.
- 1847 *Orthis filosa* J. de C. Sowerby; Davidson, p. 62, pl. 13, fig. 24.
- 1848 *Leptaena filosa* (J. de C. Sowerby); Davidson, p. 318, pl. 3, fig. 9.
- 1848 *Strophomena filosa* (J. de C. Sowerby); Phillips and Salter, p. 380.
- 1871 *Strophomena filosa* (J. de C. Sowerby); Davidson, p. 307, pl. 45, figs. 14-20.
- 1892 *Leptostrophia filosa* (J. de C. Sowerby); Hall and Clarke, p. 288.
- 1963 *Leptostrophia filosa* (J. de C. Sowerby); Holland *et al.*, pl. 3, figs. 3, 5.

Comparison. *L. (L.) filosa* differs from the common Upper Llandovery species *L. (L.) compressa* (J. de C. Sowerby) in being weakly concavo-convex to plano-convex and in having equally parvicostellate ornament, a more sharply triangular ventral muscle field with strong, relatively straight bounding ridges, and sub-parallel to weakly divergent cardinal process lobes and short socket plates. In *L. (L.) compressa* the convexity is rather variable, but is always stronger than in *filosa*; the ornament may vary from equally to unequally parvicostellate, the ventral muscle field has a broadly triangular outline and the bounding ridges are commonly weak and gently curved, and the cardinal process lobes and relatively long socket plates are widely divergent.

Description. Exterior. Very gently concavo-convex to plano-convex, commonly approaching biplanate condition; body cavity very narrow. Subquadrate to semicircular in outline, hinge-line straight, probably always forming point of maximum width, though delicate cardinal extremities frequently broken. Lateral margins of shell sub-parallel or gently curved, anterior margin rounded. Commissures smooth, anterior commissure rectimarginate.

Ventral interarea short, plane, apsacline. Delthyrium appears to be open (on material studied). Dorsal interarea plane, anacline. Notothyrium closed by convex chilidium. Ornament finely and equally parvicostellate with about 3 costellae per mm at 5-mm growth stage. Costellae low, rounded, straight along mid-line of shell but gently curved laterally. Surface of well-preserved specimens bears numerous fine, very closely spaced growth fila.

Interior of pedicle valve. Ventral process low, flat, narrow, slightly hollowed out along mid-length and produced anteriorly as slender myophragm which longitudinally bisects muscle field. Process pits narrow, elongated. Hinge-line denticulate for half its

length, denticles being borne on pair of short denticular plates. Minute hinge teeth may be present.

Muscle field sharply triangular in outline, well impressed, anteriorly flabellate, occupying about half to two-thirds of length of valve; bounded laterally by fairly strong ridges, commonly straight, but may be gently curved (medially convex). Bounding ridges arise immediately anterior to hinge, diverge antero-laterally at about 50° to one another, die out at about mid-length of muscle field, which remains unbounded anteriorly. Adductor scars narrow, elongate, separated by myophragm and extending anteriorly for just over half of length of muscle field. Diductor scars large, triangular, commonly scored by strong, radiating striae.

Area immediately lateral to muscle bounding ridges may be coarsely tuberculate and whole of internal surface is finely pseudopunctate.

Interior of brachial valve. Cardinalia supported on fairly high notothyrial platform. Cardinal process lobes narrow, plate-like, upright, sub-parallel, with attachment faces

EXPLANATION OF PLATE 57

Figs. 1–6. *Leptostrophia (Leptostrophia) filosa* (J. de C. Sowerby).

1 *a–c.* Dorsal, ventral, and lateral views of articulated shell, NMW 70.3G.7, $\times 1.3$, top 10 ft of Wenlock Limestone, old quarry on W side of road from Cwm to Ton Farm, Usk inlier, Mon., SO/3325.0180.

2. Internal moulds of 3 pedicle valves, GSM 12755, $\times 1$, Wenlock Series, Llandovery district, Carmar.; figured Davidson 1871, pl. 44, fig. 20.

3. Interior of pedicle valve, GSM 13306, $\times 1$, Wenlock Limestone, Dudley, Worcs.

4*a, b.* Artificial (calcined) internal mould of pedicle valve and counterpart exterior, NMW 70.3G.8*a, b*, $\times 1.3$, locality and horizon as for fig. 1.

5. External mould of brachial valve, lectotype, GSM Geol. Soc. Colln. 6644, $\times 1.3$, Wenlock Shale, Oldcastle, Malvern Hills.

6*a, b.* Internal mould of brachial valve and latex cast, NMW 70.3G.10, $\times 1.5$, siltstone overlying Wenlock Limestone, old quarry at Cwm, Usk inlier, Mon., SO/3331.0160.

Figs. 7, 8. *Leptostrophia (Leptostrophia) compressa* (J. de C. Sowerby).

7*a, b.* Internal mould of pedicle valve and counterpart external mould, UB 12376*a, b*, $\times 1$, base of Wenlock Series, near hedgerow, approx. 280 yd SSE of Underwood Farm, Charfield, Tortworth inlier, Glos.

8*a, b.* Internal mould of pedicle valve and counterpart external mould, CMB Cb4608*a, b*, $\times 1$, base of Wenlock Series, exposure in ditch, approx. 170 yd S.33°E. of Catherine Villa, Stone, Tortworth inlier, Glos.

Figs. 9–15. *Mclearnites coralli* sp. nov.

9*a, b.* Internal mould of pedicle valve and counterpart external mould, GSM 12753/4, $\times 1$, 'Wenlock Shale' (probably Coralliferous Series), Lindsay Bay, Pembs.

10. Internal mould of pedicle valve, holotype, GSM 12747, $\times 1$, 'Wenlock Shale' (Coralliferous Series), Slate Mill, Hasgurd, Pembs.


11. External mould of brachial valve showing interareas and part of ventral muscle field impressed by compression, GSM 12787, $\times 1$, locality and horizon as for fig. 10.

12. External mould of pedicle valve, GSM 12786, $\times 1$, locality and horizon as for fig. 10.

13. Internal mould of pedicle valve, GSM 12748, $\times 1.3$, locality and horizon as for fig. 10.

14. Internal mould of pedicle valve, SM A39419, $\times 1$, 'Calcareous Grits. ? Ludlow' (probably Sandstone Series, late Wenlock), 130 ft above base of succession, Lindsay Bay, Pembs.

15*a, b.* Internal mould of brachial valve and latex cast, NMW 70.3G.36, $\times 1$, base of Sandstone Series (probably late Wenlock), cliff section, Marloes Bay, 430 yd SW of Little Marloes Farm, Pembs., SM/7880.0710.



BASSETT, Wenlock Stropheodontidae

directed slightly postero-ventrally. Socket plates sub-parallel to cardinal process lobes or only slightly divergent.

Muscle field well impressed, truncated oval in outline, bisected longitudinally by rounded ridge which represents an anterior continuation of notothyrial platform. Muscle field bounded laterally by pair of low, broad ridges; unbounded anteriorly. Semi-oval adductor scars occupy about a quarter of width of valve. Area immediately lateral to muscle field may be coarsely tuberculate.

Dimensions of figured specimens in mm

		Length	Width
External mould of brachial valve	Lectotype, GSM Geol. Soc. Colln. 6644	21.2	29.4 est.
Interior of pedicle valve	GSM 13306	24.0	30.2 est.
Exterior of pedicle valve	GSM 13307	26.8	34.0 est.
Internal moulds of 3 pedicle valves	GSM 12755	—	—
Internal mould of pedicle valve	NMW 70.3G.9a	21.2	25.5 est.
Articulated shell	NMW 70.3G.7	17.4	—
Artificial (calcined) internal mould of pedicle valve and counterpart exterior	NMW 70.3G.8a, b	25.0	29.8
Internal mould of brachial valve	NMW 70.3G.10	—	—

Type and figured specimens. Lectotype (here selected), GSM. Geol. Soc. Colln. 6644; J. de C. Sowerby in Murchison, p. 630, pl. 13, fig. 12; from the Wenlock Shale of 'Old Castle' in the Malvern Hills, one of the localities originally mentioned by Sowerby (op. cit., p. 630). This specimen is re-figured on Plate 57, fig. 5. It was presumably one of at least two specimens available to Sowerby as he also recorded the species from Burrington in the Ludlow anticline; however, the whereabouts of any syntypes apart from the lectotype are not known.

Distribution. This species occurs fairly commonly in the upper Wenlock Shale and in the Wenlock Limestone throughout the Welsh Borderland. It is not known from early Wenlock horizons and is known only rarely in the late Wenlock mudstones of the Llandovery–Llandeilo district. No specimens have been collected from Pembrokeshire. It is also a fairly common species in the early–middle Ludlow beds of the shelf area, being recorded by Lawson (1960, p. 119) as rare or absent after Leintwardinian times.

Leptostrophia (Leptostrophia) compressa (J. de C. Sowerby)

Plate 57, figs. 7, 8

1839 *Orthis compressa* J. de C. Sowerby in Murchison, p. 638, pl. 22, fig. 12.

1967 *Leptostrophia (Leptostrophia) compressa* (J. de C. Sowerby); Cocks, p. 248, pl. 37, figs. 1–9, 11 (see also for full synonymy).

Discussion. A number of specimens of *L. (L.) compressa* have been examined by the writer from the base of the Wenlock Series in the Tortworth inlier. They were collected by Dr. M. L. K. Curtis from beds of fine sandstone which he considers to be the lateral equivalents of the basal Wenlock limestone in other parts of the inlier. Cocks (1967, pp. 248–52) described the variation in Upper Llandovery specimens of *compressa* in great detail and as the basal Wenlock specimens fall within this range there is little need to describe them here.

Cocks (1967) showed that the mean angle of divergence of the ventral muscle field of *L. (L.) compressa* increases with time in the Upper Llandovery. Insufficient material has been available to determine whether the angle of the muscle divergence of an early Wenlock population may be distinguished from that in late Llandovery populations. In two Wenlock specimens examined by the writer the muscle divergence is about 100° , but this figure falls within the range recorded by Cocks for his youngest Llandovery (C_8) population.

L. (L.) compressa has also been recorded by Cocks (1967, p. 249) from sandstone lenses in the Woolhope Limestone, of basal Wenlock age, in the May Hill inlier.

Subgenus *Leptostrophia* (*Eostropheodonta*) Bancroft 1949

Type species. *Orthis hiranensis* McCoy 1851, by original designation of Bancroft 1949, p. 9.

Discussion. The writer agrees with Cocks (1967, p. 247) that the plano-convex to gently concavo-convex subgenus *Eostropheodonta* is best regarded as a subgenus of *Leptostrophia*. Therefore, although assigned to the subfamily Stropheodontinae by Williams (1965, p. H395) *Eostropheodonta* is here removed to the Leptostrophinae. Havlíček (1967, pp. 80, 81) recognized *Eostropheodonta* as a distinct genus which he assigned to a new family Eostropheodontidae. In reaching this conclusion Havlíček (1967) suggested that *Eostropheodonta* was derived from the rafinesquinids and differs from the true stropheodontids in lacking denticulation of the cardinal margin and in lacking a ventral process which separates process pits. However, as Williams (1953a, pp. 8–10) has convincingly demonstrated, even the earliest representatives of *Eostropheodonta*, including the type species *E. hiranensis*, possess primitive denticulate plates which became modified and reduced in the phylogenetic development of the stock, resulting in the eventual disappearance of dental plates and the spread of denticles along the hinge-line; this development also took place in other stropheodontid groups and there can be little doubt that it represented a real phylogenetic trend in the family as a whole. Similarly Williams (1953a, pp. 13–15) indicated that the ventral process of stropheodontids became more pronounced with time and its weak development or absence in the earliest representatives does not justify the recognition of a separate family. Havlíček's family Eostropheodontidae is therefore not recognized here, although the writer agrees that *Eostropheodonta* itself was probably derived from a rafinesquinid ancestor. As discussed above, *Eostropheodonta* is regarded instead as an early member of the leptostrophiid stock.

Cocks (1967, p. 247) demonstrated that Llandovery specimens of *Eostropheodonta*, which possess dental plates, may occur contemporaneously with related specimens of *Leptostrophia* s.s., which lack dental plates. The description below of a Wenlock species of *Eostropheodonta* indicates that this type of coexistence continued into the Wenlock period.

Leptostrophia (*Eostropheodonta*) sp.

Plate 56, figs. 7, 8

Description. *Exterior of pedicle valve.* Valve gently convex at umbo but almost flat for remainder of length. Outline subquadrate, hinge-line straight with very short, rounded

ears. Lateral margins sub-parallel, anterior margin evenly curved. Anterior commissure rectimarginate.

Ventral interarea very weakly apsacline. Delthyrium triangular, apparently open. Fine, parallel growth-lines visible on the interarea.

Ornament unequally parvicostellate with up to 4 fine costellae occupying interspaces between low, rounded major costellae. Median costellae straight, lateral costellae gently curved. Whole surface crossed by very fine, slightly lamellose growth fila.

Interior of pedicle valve. Very short, slender dental plates diverge antero-laterally at about 90–100° to one another. Small callist occupies posterior part of delthyrial cavity. Two short, flat plates on hinge-line immediately alongside delthyrium probably represent denticular plates.

Musculature obscure but muscle field appears to have broadly triangular outline and lacks bounding ridges. Internal surface strongly impressed by external ornament.

Brachial valve unknown.

Dimensions of figured specimens in mm

		Length	Width
Internal mould of pedicle valve and counterpart external mould	NMW 70.3G.11a, b	19.3	21.0 est.
External mould of pedicle valve	NMW 70.3G.12b	20.6	—

Distribution. The above description is based on 3 poorly preserved external and internal moulds of pedicle valves from the Wenlock Shale of Craig-y-Garced in the Usk inlier, Monmouthshire (Grid. Ref. SO/3570.0275). The specimens are slightly flattened and no denticles have actually been observed because of the resulting poor preservation. However, the distinctive parvicostellate ornament and the presence of what are probably denticular plates clearly indicate the stropheodontid affinities of the specimens.

Subfamily DOUVILLININAE Caster 1939

Genus AMPHISTROPHIA Hall and Clarke 1892

Subgenus *Amphistrophia* (*Amphistrophia*) Hall and Clarke 1892

Type species. *Strophomena striata* Hall 1843, by original designation of Hall and Clarke 1892, p. 292.

The subgenus *Amphistrophia* (*Amphistrophia*) is here restricted to include only those amphistrophids which lack dental plates. A new subgenus, *Amphistrophia* (*Pembrostrophia*), is erected (see p. 325) to include forms with dental plates.

Amphistrophia (*Amphistrophia*) *whittardi* Cocks

Plate 58, figs. 2–4

1932 *Stropheodonta funiculata* (McCoy); Whittard, list facing p. 896.

1967 *Amphistrophia whittardi* Cocks, p. 261, pl. 39, figs. 3, 5, 8.

Comparison. This species closely resembles *A. (A.) funiculata* (McCoy) in outline and convexity but the two are easily distinguished by differences in musculature and ornamentation. *A. (A.) whittardi* has unequally parvicostellate radial ornament and weakly

impressed muscle fields which lack well-developed bounding ridges. In contrast *A. (A.) funiculata* has equally or subequally parvicostellate ornament and well-impressed ventral and dorsal muscle fields, both of which are bounded by strong ridges.

Description. Exterior. Semicircular to semi-elliptical in outline, about two-thirds as long as wide. Shell resupinate; pedicle valve initially gently convex, brachial valve initially gently concave. At about five-sixths of valve length both valves are geniculated smoothly and fairly gently in ventral direction. Hinge-line straight, slightly mucronate, forming point of maximum width of shell. Lateral margins evenly curved, anterior margin gently curved or prolonged slightly anteriorly to produce subtriangular outline. Anterior commissure rectimarginate.

Ventral interarea plane, apsacline. Delthyrium bears small, apical pseudodeltidium; delthyrial angle about 90°. Dorsal interarea plane, anacline. Notothyrium occupied by cardinal process.

Ornament unequally parvicostellate. Three or four fine costellae occupy broad inter-spaces between larger, rounded, major costellae.

Interior of pedicle valve. Ventral process low, flat. Dental plates absent. Pair of low denticular plates occupy median third of hinge. Muscle field weakly impressed, individual

EXPLANATION OF PLATE 58

Figs. 1a, b. *Mclearnites coralli* sp. nov. Internal and external moulds of pedicle valves showing the typical distorted preservation, NMW 70.3G.37a, b, $\times 1$, base of Sandstone Series (probably late Wenlock), cliff section, Marloes Bay, 430 yd SW of Little Marloes Farm, Pems., SM/7880.0710.

Figs. 2-4. *Amphistrophia (Amphistrophia) whittardi* Cocks.

2a, b. Internal mould of pedicle valve and counterpart external mould, NMW 70.3G.33a, b, $\times 1.5$, Wenlock mudstone, W bank of Afon Sawdde, 500 yd SE of Rhydysaint, Carms., SN/7182.2551.

3, 4. Internal moulds; 3, pedicle valve, NMW 70.3G.34, $\times 1.5$; 4, brachial valve, NMW 70.3G.35a, $\times 1.5$, locality and horizon as for fig. 2.

Figs. 5-16. *Amphistrophia (Amphistrophia) funiculata* (McCoy).

5a-d. Dorsal, ventral, lateral, and posterior views of complete shell, NMW 70.3G.13, $\times 1.5$, Wenlock Limestone, old quarry and tip on W side of Lincoln Hill, Ironbridge, Salop., SJ/6695.0381.

6, 7, 8. Brachial valves; 6, exterior, NMW 70.3G.14, $\times 1.5$; 7, interior, NMW 70.3G.15, $\times 2$; 8, exterior, NMW 70.3G.16, $\times 2$; locality and horizon as for fig. 5.

9. External mould of pedicle valve, lectotype, NMI 111 11, $\times 1.5$, Silurian, Doonquin, Dingle, Co. Kerry, Ireland.

10, 11. Interiors of pedicle valves; 10, NMW 70.3G.17, $\times 1.5$; 11, NMW 70.3G.18, $\times 1.5$, locality and horizon as for fig. 5.

12a, b. Internal mould of brachial valve and latex cast, NMW 70.3G.19, $\times 1.5$, siltstone overlying Wenlock Limestone, old quarry at Cwm, Usk inlier, SO/3331.0160.

13a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.20, $\times 1.5$, Wenlock Shale, temporary drainage trench from Cwm to railway line at Monkswood, Usk inlier, Mon., SO/3335.0173 to 3371.0215.


14a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.21a, $\times 1.5$, locality and horizon as for fig. 12.

15a, b. Internal mould of brachial valve and latex cast, NMW 70.3G.22a, $\times 1.5$, locality and horizon as for fig. 12.

16a-d. Internal mould of brachial valve and latex cast, counterpart external mould and latex cast, NMW 70.3G.23a, b, $\times 1.5$, locality and horizon as for fig. 13.



1a




1b



2a



2b



3



4



5a




5d



5b



5c



6



7



8



9




10



11



12a



12b



13a




13b



14a



14b




15a



15b



16a



16b



16c



16d

BASSETT, Wenlock Stropheodontidae

scars not discernible. Very faint myophragm bisects muscle field longitudinally. Extremely short, faint ridges arise near umbo and diverge antero-laterally at about 90° to one another to bound muscle field postero-laterally; these ridges die out rapidly and greater proportion of muscle field remains unbounded.

Interior of brachial valve. Notothyrial platform very low. Cardinal process lobes sub-parallel, postero-ventrally directed. Sockets shallow, elongated; socket plates low, fairly wide, antero-laterally divergent at about 120° to one another. Muscle field not impressed, bounding ridges absent.

Dimensions of figured specimens in mm

		Length	Width
Internal mould of pedicle valve and counterpart external mould	NMW 70.3G.33a, b	11.2	18.6
Internal mould of pedicle valve	NMW 70.3G.34	11.9	16.7
Internal mould of brachial valve	NMW 70.3G.35a	11.7	—

Type and figured specimens. The specimens of *A. (A.) whittardi* figured by Cocks (1967, pl. 39, figs. 3, 5, 8), including the holotype, are in the Oxford University Museum (see Cocks, 1967, p. 262).

Distribution. This species has been collected by the writer from a single locality in the Sawdde gorge section, Carmarthenshire (Grid. Ref. SN/7176.2562) in beds mapped by Williams (1953b, p. 198) as lower Wenlock in age. It was previously recorded by Cocks (1967, p. 263) only from the late Llandovery Purple Shales of the Welsh Borderland, but has also been examined by the writer in collections, now housed in the City Museum, Bristol, made by Dr. M. L. K. Curtis from the Upper Llandovery of the Tortworth inlier.

Amphistrophia (Amphistrophia) funiculata (McCoy)

Plate 58, figs. 5–16

- 1846 *Orthis funiculata* McCoy, p. 30, pl. 3, fig. 11.
- 1847 *Leptaena funiculata* (McCoy); Davidson, p. 57, pl. 12, figs. 5–8.
- 1848 *Leptaena funiculata* (McCoy); Davidson, p. 317, pl. 3, fig. 5.
- 1852 *Leptaena (Strophomena) funiculata* (McCoy); McCoy, p. 244.
- 1861 *Strophomena funiculata* (McCoy); Lindström, p. 372.
- 1871 *Strophomena funiculata* (McCoy); Davidson, p. 290, pl. 40, figs. 9–13.
- 1963 *Strophonella funiculata* (McCoy); Holland *et al.*, pl. 3, figs. 5, 6, 7.
- ? 1967 *Amphistrophia funiculata* (McCoy); Havlíček, p. 186, pl. 29, figs. 1, 2, 4, 5, 9.

Discussion. McCoy's (1846, pl. 3, fig. 11) original specimen of '*Orthis*' *funiculata*, from the Silurian of Doonquin, Dingle, Co. Kerry, Ireland, is an extremely poorly preserved external mould of a pedicle valve. The antero-lateral margins of this shell have a rather angular outline in comparison with the evenly curved margins of specimens from the Welsh Borderland and neighbouring areas which are commonly identified as *funiculata*. Unfortunately material from Dingle is very rare so it is not possible to judge at this stage whether the difference in shape between the lectotype and Anglo-Welsh shells is taxonomically significant. However, three poorly preserved specimens of *funiculata*, collected from Dingle by Professor C. H. Holland, show some variation in outline and

have ventral muscle fields which are morphologically inseparable from those in Anglo-Welsh specimens. It seems advisable, therefore, to retain the name *funiculata* for the latter. If future studies should indicate that populations from Dingle have a significantly different shape, the differences will probably be best expressed by splitting off the Anglo-Welsh shells at no more than the subspecies level.

Comparison. A comparison of this species with *A. (A.) whittardi* is given earlier (p. 319). *A. (A.) funiculata* differs from *A. (A.) striata* (Hall), the type species of *Amphistrophia* from the Waldron Shale of Indiana, in having a transversely semicircular outline, a relatively longer ventral muscle field which may be completely bounded by ridges, and in being fairly sharply geniculate. *A. (A.) striata* has a subquadrate outline, is only gently resupinate, and has a ventral muscle field which is confined to the posterior half of the valve and is unbounded anteriorly.

Description. Exterior. Outline semicircular to semi-elliptical, commonly about two-thirds to half as long as wide. Maximum width at straight hinge-line, which may be produced as short rounded ears. Shells resupinate; pedicle valve initially gently convex, brachial valve initially gently concave. Valves are then geniculated fairly sharply ventrally, commonly between 10- and 15-mm growth stages. Commissures crenulate, anterior commissure rectimarginate. Ventral interarea plane, apsacline, about 3 times as long as dorsal interarea which is plane, anacline. Delthyrium bears small, convex, apical pseudodeltidium and is partially occupied by cardinal process. Delthyrial angle about 80°. Notothyrium filled by cardinal process, which is flanked by pair of slender chilidial plates. Impress of internal denticulation may be visible on both interareas. Ornament equally to sub-equally parvicostellate with 4-5 low, rounded costellae per mm at 10-mm growth stage. New costellae arise both by insertion and bifurcation.

Interior of pedicle valve. Ventral process low, produced anteriorly as slender myophragm which longitudinally bisects muscle field. Process pits shallow. Median third of hinge denticulate, denticles being borne on pair of well-defined denticular plates.

Muscle field strongly impressed, longitudinally oval in outline, occupying from half to four-fifths of total length of valve. Strong, medially concave ridges bound muscle field laterally, and may die out at point where muscle field reaches maximum width in its anterior half; muscle field may then remain unbounded anteriorly; in some specimens bounding ridges completely enclose muscle field, intersecting myophragm at its anterior extremity. Adductor scars narrow, lanceolate, confined to posterior half of muscle field and bounded laterally and anteriorly by large, oval diductor scars.

Internal surface may be coarsely tuberculate and may show impress of external ornament.

Interior of brachial valve. Cardinal process lobes postero-ventrally directed, U-shaped in cross-section, pear-shaped in outline, separated medially by rounded groove. Sockets fairly deep, bounded anteriorly by strong socket plates diverging antero-laterally at about 100-120° to one another and which may be crenulated on their posterior edges.

Muscle field well impressed, truncated oval in outline, occupying about one-quarter of length of valve. Adductor scars bounded laterally by strong, medially concave ridges;

unbounded anteriorly. Muscle scars separated longitudinally by low, rounded myophragm arising below the cardinalia and tapering anteriorly.

Area immediately lateral to muscle field commonly coarsely tuberculate.

Dimensions of figured specimens in mm

		<i>Length</i>	<i>Width</i>
External mould of pedicle valve	Lectotype, NMI 111 11	18.2	27.0 est.
Complete shell	NMW 70.3G.13	12.6	19.5 est.
Exterior of pedicle valve	NMW 70.3G.14	13.0	20.0
Interior of brachial valve	NMW 70.3G.15	—	—
Exterior of pedicle valve	NMW 70.3G.16	11.6	18.5
Interior of pedicle valve	NMW 70.3G.17	15.1	19.6 est.
Interior of pedicle valve	NMW 70.3G.18	12.7	20.1
Internal mould of brachial valve	NMW 70.3G.19	11.6	19.0 est.
Internal mould of pedicle valve	NMW 70.3G.20	12.8	24.0 est.
Internal mould of pedicle valve	NMW 70.3G.21a	14.0	23.6 est.
Internal mould of brachial valve	NMW 70.3G.22a	13.6	21.5
Internal mould of brachial valve and counterpart external mould	NMW 70.3G.23a, b	13.4	23.3

Type specimens. Lectotype (here selected, figured McCoy 1846, pl. 3, fig. 11), NMI 111 11. It is refigured on Plate 58, fig. 9. This is one of 2 specimens that were available to McCoy from the collection of Sir Richard Griffith; the other specimen is also in the National Museum of Ireland but is unregistered.

Distribution. *A. (A.) funiculata* occurs very commonly in the upper Wenlock Shale and the Wenlock Limestone throughout the Welsh Borderland and less commonly in the late Wenlock sandstones and mudstones of Tortworth and South Wales. It is not known from Pembrokeshire and does not occur in beds of early Wenlock age. The species ranges through into the Ludlovian where it is recorded by Lawson (1960, p. 117) as being rare or absent after Bringewoodian times. On Gotland it is recorded by Hede (1960) from localities in the Slite and Mulde Marls; both of these divisions are considered to be of Wenlock age.

Amphistrophia (Amphistrophia) cf. euglyphoides Hortedahl

Plate 59, figs. 1-4

cf. 1916 *Amphistrophia euglyphoides* Hortedahl, p. 65, pl. 6, figs. 7-9.

Comparison. A few poorly preserved specimens examined by the writer from the base of the Wenlock succession in the Tortworth inlier closely resemble *A. (A.) euglyphoides* Hortedahl in details of shape, convexity, ornament, and configuration of the ventral muscle field. However, until well-preserved specimens, including dorsal interiors, are collected from Tortworth, the comparison with *euglyphoides* must remain tentative. The pedicle valves of *A. (A.) euglyphoides* figured by Hortedahl (1916, pl. 6, fig. 8) are refigured for comparison on Plate 59, fig. 5.

A comparison of *A. (A.) cf. euglyphoides* and *A. (P.) freshwaterensis* is given on p. 326.

Description. Exterior. Shell resupinate, gently concavo-convex initially, becoming gently convexi-concave after length of about 10 mm. Outline subquadrate, hinge-line

straight, cardinal angles rounded. Lateral margins sub-parallel, straight to gently curved, anterior margin evenly rounded.

Ventral interarea apsacline, dorsal interarea anacline. Nature of delthyrium and notothyrium not observed.

Radial ornament fine, equally parvicostellate with approximately 4–5 low, rounded costellae per mm at 5-mm growth stage.

Interior of pedicle valve. Ventral process flat, produced anteriorly as very slender myophragm. Muscle field sub-oval, confined to posterior half of valve and occupying about one-third of valve width. Individual scars not discernible. Low, gently curved ridges bound muscle field laterally, but unbounded anteriorly where margin may be slightly flabellate.

Interior of brachial valve. Unknown.

Dimensions of figured specimens in mm

		Length	Width
Interior of pedicle valve	GSM DEW9543A	22.5	28.0 est.
Exterior of brachial valve	GSM DEW9543B	16.1	19.1 est.
Interior of pedicle valve	GSM DEW9548	—	—
Exterior of brachial valve	CMB Cb4610	15.0	17.9

Distribution. *A (A.)* cf. *euglyphoides* is known only from the Tortworth inlier where it occurs rarely in the limestone band at the base of the Wenlock Series and in sandstone

EXPLANATION OF PLATE 59

Figs. 1–4. *Amphistrophia (Amphistrophia)* cf. *euglyphoides* Høltedahl.

1. Exterior of brachial valve, CMB Cb4610, $\times 1.3$, Wenlock Series, Whitfield, Tortworth inlier, Glos.
- 2, 4. Interiors of pedicle valves; 2, GSM DEW9548, $\times 1.3$; 4, GSM DEW9543A, $\times 1.5$; basal Wenlock limestone, Rifle Cottage Quarry, Tortworth inlier, Glos.
3. Exterior of brachial valve, GSM DEW9543B, $\times 1.3$, locality and horizon as for fig. 2.

Figs. 5–6. *Amphistrophia (Amphistrophia)* *euglyphoides* Høltedahl.

5. Interiors of pedicle valves, PMO L0116, $\times 1$, Stage 9c, Langø, off Holmestrand, Oslo district, Norway; figured Høltedahl 1916, pl. 6, fig. 8.
6. Exteriors of pedicle valves, PMO L0122, $\times 1$, locality and horizon as for fig. 5; figured Høltedahl 1916, pl. 6, fig. 7.

Fig. 7. *Amphistrophia (Amphistrophia)* sp. Internal mould of pedicle valve, GSM Hc992, XI, late Wenlock mudstone, old overflow trench near Middleton Hall, Carms., SN/5265.1879.

Fig. 8. *Amphistrophia* sp. Internal mould of pedicle valve, NMW 70.3G.47, $\times 1$, base of Sandstone Series (probably late Wenlock), cliff section above Marloes Bay, approx. 750 yd NW of Little Marloes Farm, Pembs., approx. SM/7835.0765.

Figs. 9–14. *Amphistrophia (Pembrostrophia)* *freshwaterensis* subgen. et sp. nov. Top of Wenlock (of Dixon 1921), large reef on beach approx. 30 ft above lowest beds, S side of Freshwater East Bay, Pembs., SS/0165.9753.

- 9a, b. Internal mould of pedicle valve and latex cast, holotype, NMW 70.3G.25, $\times 1.25$.
- 10a, b. External mould of brachial valve and latex cast, NMW 70.3G.26, $\times 1$.
11. Internal mould of brachial valve, NMW 70.3G.27, $\times 1.5$.
- 12 a–c. Internal mould of brachial valve, latex cast, and counterpart external mould, NMW 70.3G.28a, b; a, $\times 1$; b, c, $\times 1.25$.
- 13a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.29, $\times 1.25$.
- 14a, b. Internal mould of brachial valve and latex cast, NMW 70.3G.30, $\times 1.25$.




1



2



3



4




5




6




8




7




9a



9b



10a




10b




11



12a



12b




12c



13a



13b



14a



14b

BASSETT, Wenlock Stropheodontidae

beds which are considered by Dr. M. L. K. Curtis to be lateral equivalents of the basal limestone in some parts of the inlier.

Amphistrophia (*Amphistrophia*) sp.

Plate 59, fig. 7

A single specimen of *Amphistrophia* in the collections of the Geological Survey Museum (GSM Hc992), from the late Wenlock of Middleton Park, Carmarthenshire, differs from any other specimen described here mainly in its large size and longitudinally elongated outline. Associated with the specimen is a brachial valve which may belong to the same species. A full description of this species is deferred until more material can be collected to allow its relationships with other species to be assessed.

Dimensions in mm. Length, 41.7; width 46.0 est.

Subgenus *Amphistrophia* (*Pembrostrophia*) subgen. nov.

Type species. *Amphistrophia* (*Pembrostrophia*) *freshwaterensis* sp. nov.

Diagnosis. A subgenus of *Amphistrophia* possessing dental plates.

Discussion. In the late Wenlock/Ludlow rocks of Freshwater East Bay, Pembrokeshire, there occurs a fairly large gently resupinate amphistropheiid which possesses small but distinctive dental plates and which thus morphologically represents the 'Eo' stage (Williams 1953a, p. 30) of *Amphistrophia* s.s. As the earliest species of *Amphistrophia* yet recorded is *A. (A.) whittardi* from the late Llandovery of the Welsh Borderland (Cocks 1967, p. 262) one would normally expect 'Eo' forms of the genus to occur in beds of pre-Upper Llandovery age. The presence of amphistropheiiids with dental plates in Wenlock/Ludlow rocks is therefore somewhat surprising. If it is assumed that *Amphistrophia* followed the normal pattern of stropheodontid evolution, and that forms with dental plates ('Eo' stage) gave rise to forms without dental plates ('Meso' stage), the Pembrokeshire specimens described here may be considered to represent conservative members of an ancestral amphistropheiid stock existing contemporaneously with descendant forms. In other respects, however, *A. (P.) freshwaterensis* appears to represent a fairly well-advanced species since it has patterns of ribbing and musculature more closely comparable with those in *A. (A.) funiculata* than in the early species *A. (A.) whittardi*; there is a possibility, therefore, that *A. (P.) freshwaterensis* developed from a stock without dental plates and that these structures were introduced as a secondary phenomenon, perhaps in response to local environmental conditions.

It is interesting to note that within one population of *A. (P.) freshwaterensis* the development of dental plates is rather variable and in a few specimens they are represented instead by low flat pads of shell material (e.g. Pl. 59, fig. 13). If the presence of plates is explained purely in evolutionary terms then it is probable that the species is approaching the stage where plates will be completely lost. Some support for this alternative is provided by a few specimens of *Amphistrophia* collected from low in the Sandstone Series of Marloes Bay in Pembrokeshire; these specimens are probably slightly younger

than those from Freshwater East, but are from a similar lithology, and yet all lack dental plates (e.g. Pl. 59, fig. 8). The lack of detailed knowledge of the stratigraphy of the Silurian in Pembrokeshire precludes meaningful discussion on the evolutionary position of *A. (Pembrostrophia)*, and it is clear that its relationship will not be evaluated until undoubtedly older and younger populations from the same area can be studied in detail. However, the presence of well-defined dental plates in the majority of the specimens in a population of *A. (P.) freshwaterensis* certainly warrants subgeneric status. Since these specimens with dental plates post-date the earliest (Llandovery) species of *Amphistrophia* (*Amphistrophia*), the term 'Eo', which implies a time relationship, is not used as a prefix to the generic name as is normally the practice in stropheodontid systematics. Instead, the subgeneric name *Pembrostrophia* is here proposed.

Amphistrophia (Pembrostrophia) freshwaterensis sp. nov.

Plate 59, figs. 9–14; Plate 60, figs. 1, 2

Diagnosis. Subquadrate to sub-circular *Amphistrophia*, gently resupinate, dental plates small, variably developed. Ornament finely and equally parvicostellate. Ventral and dorsal muscle fields well impressed, with well-developed bounding ridges.

Comparison. No other species of *A. (Pembrostrophia)* are yet known. *A. (P.) freshwaterensis* closely resembles *A. (A.) euglyphoides* Høltedahl in details of shape, convexity, musculature, and ornament. However, in *euglyphoides* (and in specimens described herein as cf. *euglyphoides*) true dental plates are never seen although low pads of calcite may be present, similar to those in end members of populations of *freshwaterensis*. The two species are undoubtedly very closely related, but until the age of the Pembrokeshire sections can be accurately determined it is not possible to assess their relationships in amphistrophiid phylogeny.

Description. Exterior. Subquadrate to sub-circular in outline with point of maximum width at or slightly posterior to mid-length. Hinge-line straight, cardinal angles rounded; hinge width equal to about nine-tenths of maximum width. Shell resupinate; pedicle valve initially gently convex, becoming gently concave anteriorly; brachial valve initially plane or gently concave, becoming gently convex anteriorly, sometimes tending to flatten out near anterior margin. Lateral margins sub-parallel or gently curved, anterior margin evenly rounded or sub-parallel to hinge. Anterior commissure rectimarginate.

Ventral interarea plane, strongly apsacline. Delthyrial angle about 110°. Delthyrium bears small, gently convex, apically situated pseudodeltidium. Dorsal interarea short, plane, anacline. Nature of notothyrium not observed.

Radial ornament finely and equally parvicostellate. Costellae low, rounded, with 4–5 per mm at 5-mm growth stage of pedicle valve. Fine, irregularly spaced concentric growth-lines visible on well-preserved specimens.

Interior of pedicle valve. Pseudodeltidium supported by low ventral process, concave along its mid-length and produced anteriorly as thin myophragm of constant width. Process pits shallow. Hinge-line denticulate for about one-quarter of its length, denticles

being borne on pair of well developed denticular plates. At their junction with hinge-line, bases of denticular plates supported by low, flat dental plates. In some specimens dental plates almost totally reduced to low pad of shell material between muscle bounding ridges and hinge-line.

Muscle field well impressed, sub-oval in outline, occupying up to half valve length and up to half width. Strong, medially concave ridges bound muscle field laterally, may die out in anterior half of muscle field to leave scars unbounded anteriorly, or may curve medially and then posteriorly to intersect myophragm and completely enclose muscle field. Adductor scars narrow, lanceolate, tapering anteriorly, commonly confined to posterior half of muscle field. Diductor scars large, oval, enclosing adductor scars laterally and anteriorly. Anterior margin of muscle field may be flabellate and whole muscle area may be scored by faint, radial striae.

Internal surface of valve may be faintly impressed by external ornament.

Interior of brachial valve. Cardinalia supported on high notothyrial platform. Cardinal process lobes narrow, pear-shaped, almost meeting posteriorly, separated by rounded groove which may bear narrow ridge in its posterior half. Attachment faces of lobes grooved longitudinally and directed postero-ventrally. Socket ridges long, antero-laterally divergent at about 120° to one another, arising slightly lateral to posterior tips of cardinal process lobes and extending almost to lateral margins of muscle field. Sockets fairly deep.

Muscle field well impressed, occupying about one-quarter of length of valve, unbounded anteriorly; bounded laterally by broad, strong ridges, and bisected longitudinally by high, convex median ridge which tapers anteriorly and extends to anterior margin of subrectangular scars.

Denticles present on hinge, complementary to those of pedicle valve.

Dimensions of figured specimens in mm

		<i>Length</i>	<i>Width</i>
Internal mould of pedicle valve	Holotype, NMW 70.3G.25	21.6	25.0
External mould of pedicle valve	NMW 70.3G.26	20.2	25.4
Internal mould of brachial valve	NMW 70.3G.27	—	—
Internal mould of pedicle valve and counterpart of external mould	NMW 70.3G.28a, b	21.0	26.4 est.
Internal mould of pedicle valve	NMW 70.3G.29	21.1	26.6 est.
Internal mould of brachial valve	NMW 70.3G.30	19.8	—
Internal mould of pedicle valve	NMW 70.3G.31	17.5 est.	24.0 est.
Internal mould of pedicle valve	NMW 70.3G.32	—	—

Holotype. Internal mould of pedicle valve, NMW 70.3G.25, figured Plate 59, fig. 9.

Distribution. This species is known at present only from Freshwater East Bay in Pembrokeshire. It is very common on the south side of the bay in beds considered by Dixon (1921, p. 13) to be at the top of the Wenlock Series but which are considered by the writer to be of late Wenlock or Ludlow age. It is also present in collections made by Dr. V. G. Walmsley from slightly younger beds (Ludlow of Dixon 1921, p. 14) on both the north and south sides of the bay.

Genus MCLEARNITES Caster 1945 [pro *Mclearnia* Caster 1939]

Type species. *Brachyprion mertonii* McLearn 1928, by original designation of Caster 1939, p. 28.

Discussion. The genus *Mclearnites* was erected [pro *Mclearnia*] by Caster (1939, 1945) with *Brachyprion mertonii* McLearn, from the Stonehouse Formation of Nova Scotia, as its type species. Little or no mention of the genus was then made until Williams (1953a, p. 47; 1965, p. H402) placed it in synonymy with *Shaleria*, within the subfamily Shaleriinae. Re-examination by the writer of a small collection of topotype specimens of *B. mertonii*, collected by Dr. V. G. Walmsley, suggests, however, that *Mclearnites* is best assigned to the Douvillinae and is generically distinct from *Shaleria*. *Shaleria* is characterized by having long, narrow, sub-parallel ventral muscle scars which are commonly bounded completely by ridges, whereas *Mclearnites* has a triangular-shaped

EXPLANATION OF PLATE 60


- Figs. 1, 2. *Amphistrophia* (*Pembrostrophia*) *freshwaterensis* subgen. et sp. nov. Large reef on beach approx. 30 ft. above lowest beds, S side of Freshwater East Bay, Pembs., SS/0165.9753.
- 1a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G.31, $\times 1$, top of Wenlock (of Dixon 1921).
- 2a, b. Internal mould of pedicle valve and latex cast, NMW 70.3G. 32, $\times 1$.
- Fig. 3. *Pholidostrophia* (*Mesopholidostrophia*) *laevigata* (J. de C. Sowerby), holotype, GSM Geol. Soc. Colln. 6635, $\times 2$, Wenlock Shale, Burrington, near Ludlow, Salop.
- Figs. 4, 5. *Pholidostrophia* (*Mesopholidostrophia*) cf. *laevigata* (J. de C. Sowerby).
4. Exterior of pedicle valve, SM A16544, $\times 2$, Lower Ludlow Shales, Woolhope inlier, Herefordshire.
5. Exterior of pedicle valve, NMW 70.3G.46, $\times 2$, Wenlock Limestone, quarry on N side of Ridgeway, $\frac{1}{4}$ mile SE of Ockeridge Farm, Malvern Hills, SO/7532.3967.
- Figs. 6–9. *Pholidostrophia* (*Mesopholidostrophia*) *salopiensis* Cocks. Wenlock mudstone, W bank of Afon Sawdde, 500 yd. SE of Rhydsaint, Carms., SN/7182.2551.
- 6a, b. Ventral and postero-ventral views of internal mould of pedicle valve, NMW 70.3G.38, $\times 2$.
7. Internal mould of pedicle valve, NMW 70.3G.39a, $\times 2$.
- 8a, b. Internal mould of pedicle valve and counterpart external mould, NMW 70.3G.40a, b, $\times 2$.
9. External mould of brachial valve, NMW 70.3G.41a, $\times 2$.
- Figs. 10–13. *Pholidostrophia* (*Mesopholidostrophia*) *deflecta* sp. nov.
- 10a, b. Partially exfoliated exterior of pedicle valve and counterpart impression, holotype, NMW 70.3G.42a, b, $\times 1.5$, top 10 ft of Wenlock Limestone, old quarry on W side of road from Cwm to Ton Farm, Usk inlier, Mon., SO/3325.0180.
11. Artificial (calcined) internal mould of pedicle valve, NMW 70.3G.43a, $\times 1.5$, locality and horizon as for fig. 10.
12. Internal mould of pedicle valve, NMW 70.3G.44, $\times 1.5$, siltstone above Wenlock Limestone, NW corner of old quarry behind Glascoed-fach Farm, Usk inlier, Mon., SO/3291.0210.
- 13a, b. Internal mould of brachial valve and latex cast, NMW 70.3G.45, $\times 1.5$, locality and horizon as for fig. 12.
- Figs. 14–16. *Pholidostrophia* (*Mesopholidostrophia*) *fletcheri* (Davidson). Wenlock Limestone, Benthall Edge, Salop.
14. Exterior of brachial valve, BM BB32804 (ex B5821), $\times 1.5$.
- 15a–c. Ventral, dorsal and lateral views of articulated shell, lectotype, BM B5821, $\times 1.5$.
16. Exterior of brachial valve, BM BB32803 (ex B5821), $\times 1.5$.



1a



1b



2a



2b



3



4



5



8a



6a



6b



7



8b



9



10a



10b



11



12



13a



13b



14



15a



15b



15c



16

BASSETT, Wenlock Stropheodontidae

ventral muscle field with low, curved lateral bounding ridges which are typical of the douvillinids. The pedicle valve of *Shaleria* is also commonly plane whereas in *Mclearnites* it is gently to fairly strongly convex. The recognition of *Mclearnites* as a distinct genus has also been suggested recently by Boucot *et al.* (1966, addendum, p. 41) and Cocks (1967, p. 251). *Mclearnites* may be a senior synonym of *Mesodouvillina* Williams 1950 (Boucot *et al.* 1966, addendum, p. 41) but little comment can be made here in the absence of comparative material.

Mclearnites coralli sp. nov.

Plate 57, figs. 9–15; Plate 58, fig. 1

1967 *Mclearnites* sp. Cocks, pl. 37, fig. 10.

Diagnosis. Very gently concavo-convex to plano-convex *Mclearnites*. Ventral muscle field broadly triangular with gently curved lateral margins defined by low bounding ridges. Ventral myophragm weak. Socket ridges weak. Dorsal muscle field faintly impressed, lacking bounding ridges. Ornament fine, sub-equally parvicostellate.

Comparison. *Mclearnites coralli* differs from *M. mertonii* in having slightly finer radial ornament, a wider ventral muscle field which is bisected by a very faint myophragm, and a weakly impressed dorsal muscle field which lacks lateral bounding ridges. *M. mertonii* has a relatively narrow, triangular ventral muscle field which is bisected by a well-developed myophragm, and a strongly impressed dorsal muscle field which is bounded laterally by a pair of well-developed ridges.

Description. Exterior. Outline subquadrate to semicircular, pedicle valve gently to moderately convex, brachial valve plane to gently concave. Hinge-line straight, forming point of maximum width of shell, cardinal angles rounded or slightly mucronate. Lateral margins of shell almost straight, sub-parallel; anterior margin evenly curved, anterior commissure rectimarginate.

Ventral interarea plane, weakly apsacline. Delthyrium appears to be open (on material available), with delthyrial angle varying from about 80–110°. Notothyrium appears to be occupied by cardinal process but dorsal interarea not studied in detail.

Ornament equally to sub-equally parvicostellate, with about 4 costellae per mm over whole surface of shell. Costellae low, rounded, increasing mainly by insertion. Very fine, closely spaced growth fila visible on well-preserved specimens.

Interior of pedicle valve. Teeth obsolete, dental plates absent. Hinge-line denticulate along its median third, denticles being borne on pair of low denticular plates.

Muscle field broadly triangular in outline, confined to posterior half of valve and occupying about one-third valve width. Laterally muscle field bounded by pair of gently but distinctly curved ridges, medially concave. Faint myophragm bisects muscle field longitudinally. Adductor scars narrow, lanceolate, situated adjacent to myophragm and bounded laterally by broad diductor scars.

Extra-muscular area finely pseudopunctate, showing impress of external ornament.

Interior of brachial valve. Cardinalia raised on low notothyrial platform produced anteriorly as very low convex median ridge. Cardinal process lobes strong, slightly expanded anteriorly, U-shaped in cross-section and divergent antero-laterally at about 90° to one another. Sockets shallow, socket ridges low, broad, antero-laterally divergent at about 120–130° to one another.

Muscle field weakly impressed, longitudinally bisected by median ridge. Truncated oval muscle scars unbounded laterally and anteriorly.

Internal surface of valve lightly impressed by external ornament.

Dimensions of figured specimens in mm

All of the specimens are slightly distorted.

		Length	Width
Internal mould of pedicle valve	Holotype, GSM 12747	20.9	25.8
Internal mould of pedicle valve and counterpart external mould	GSM 12753/4	25.8	23.0
External mould of brachial valve	GSM 12787	29.1	—
External mould of pedicle valve	GSM 12786	28.4	36.0 est.
Internal mould of pedicle valve	GSM 12748	13.0	16.4 est.
Internal mould of pedicle valve	SM A39419	—	—
Internal mould of brachial valve	NMW 70.3G.36	17.7	—

Holotype. Internal mould of pedicle valve, GSM 12747, from the 'Wenlock Shale' of Slate Mill, Hasguard, Pembrokeshire. The exact locality is unknown but the outcrops around Slate Mill are in the Coralliferous Series (see Dixon 1921) and are probably of late Wenlock age.

Distribution. *M. coralli* is known only from the Silurian of south Pembrokeshire. It has been collected by the writer from high in the Coralliferous Series at the reservoir at Winsle (SN/8333.0929) and from the cliff section at Marloes Bay, and from low in the Sandstone Series at the latter locality. Further material examined in the Geological Survey Museum and the Sedgwick Museum is from the Coralliferous Series and/or low in the Sandstone Series of Slate Mill and Lindsway Bay. All these horizons are probably of late Wenlock age.

Subfamily PHOLIDOSTROPHIINAE Stainbrook 1943

Genus PHOLIDOSTROPHIA Hall and Clarke 1892

Subgenus *Pholidostrophia* (*Mesopholidostrophia*) Williams 1950

Type species. See discussion below.

Discussion. *Pholidostrophia* (*Mesopholidostrophia*) was erected by Williams (1950, p. 278) with *P. (M.) nitens* Williams 1950 as its type species. Examination of topotype specimens of *nitens*, collected from Gotland by the writer, reveals that this species is close to and possibly conspecific with the shell described by J. de C. Sowerby as *Leptaena laevigata* (in Murchison 1839, p. 629, pl. 13, fig. 3). Unfortunately only a single specimen of *laevigata* (the holotype) is known at present and attempts to collect topotype material from the Wenlock Shale at Burrington, Shropshire, have so far proved to be unsuccessful; it is therefore not possible at this stage to compare *laevigata* and *nitens* in detail. Populations of *nitens* tend to show stronger ventral curvature than the holotype of *laevigata*, though individual specimens may be closely comparable, and it

remains to be seen whether a population of *laevigata* would exhibit the same variation. Should this prove to be so, then the writer considers that *nitens* should be regarded as a junior synonym of *laevigata*, which would thus become the type species of *Pholidostrophia* (*Mesopholidostrophia*).

Lindström (1861, p. 374) also realized the relationship of *laevigata* with *nitens* when he stated that *L. margaritacea* Angelin and *laevigata* were possibly identical. Examination of Angelin's collection of *L. margaritacea* (SMNH Br32147-58) reveals that the shells are identical with *P. (M.) nitens*; however, the name *margaritacea* must be regarded as a *nomen nudum* since Lindström's reference is to a manuscript name and neither he nor Angelin described or figured the species. Recently Harper, Johnson, and Boucot (1967, p. 415) showed that *P. (M.) nitens* is a senior synonym of *Lissoostrophia* (*Mesolissoostrophia*) *pellucida* Williams 1950.

Pholidostrophia (*Mesopholidostrophia*) *laevigata* (J. de C. Sowerby)

Plate 60, fig. 3

- 1839 *Leptaena laevigata* J. de C. Sowerby in Murchison, p. 629, pl. 13, fig. 3.
- 1847 *Leptaena laevigata* J. de C. Sowerby; Davidson, p. 59, pl. 12, fig. 30.
- 1848 *Leptaena laevigata* J. de C. Sowerby; Davidson, p. 319.
- 1871 *Leptaena* (?) *laevigata* J. de C. Sowerby; Davidson, p. 328, pl. 49, figs. 1, 2, *non* figs. 3-12.
- 1967 *Pholidostrophia* (*Mesopholidostrophia*) *laevigata* (J. de C. Sowerby); Harper *et al.*, p. 414.

Description of holotype. Exterior of pedicle valve. Valve small, gently convex, semi-circular in outline. Hinge-line straight, produced as delicate mucronate ears. Lateral and anterior margins evenly curved. Umbo low.

Shell substance slightly nacreous. Extremely faint, rounded, widely spaced ribs visible beneath nacreous shell. Surface covered with very fine, closely spaced growth fila.

Dimensions in mm. Hinge width, 9.3; length 5.4.

Holotype (by monotypy), J. de C. Sowerby in Murchison 1839, pl. 13, fig. 3, in the Geological Survey Museum (Geol. Soc. Colln. 6635), from the Wenlock Shale of Burrington, near Ludlow, Shropshire. It is refigured on Plate 60, fig. 3. Poorly preserved specimens collected by the writer from the Wenlock Limestone of the Ridgeway, $\frac{1}{4}$ mile SE of Ockeridge Farm in the Malvern Hills (Grid. Ref. SO/3967.7532) possibly also belong to *P. (M.) laevigata*.

Pholidostrophia (*Mesopholidostrophia*) *deflecta* sp. nov.

Plate 60, figs. 10-13

Diagnosis. Relatively large *Pholidostrophia* (*Mesopholidostrophia*) in which the periphery of the pedicle valve is gently deflected dorsally.

Comparison. In size and outline *P. (M.) deflecta* most closely resembles *P. (M.) fletcheri* (Davidson) but it is easily distinguished from this and all other pholidostrophiids by the characteristic dorsal deflection in the pedicle valve.

Description. Exterior. Shells relatively large for genus, plano-convex to gently concavo-convex, with curvature of pedicle valve slightly greater. Periphery of pedicle valve shows very gentle but well-defined dorsal lip or deflection. Hinge-line straight, forming point of maximum width of shell and produced as short, rounded ears. Margins of shell smoothly curved.

Ventral interarea plane, weakly apsacline, up to 3 times as long as dorsal interarea, which is plane, anacline. Delthyrium may bear very small apical pseudodeltidium but in many specimens appears to be open; delthyrial angle about 60°. Notothyrium occupied by cardinal process.

Surface of shell nacreous, lacking well-defined radial ornament. However, outer, nacreous shell layer commonly not preserved and in these cases fine parvicostellate ornament is visible. Very fine, concentric growth fila present.

Interior of pedicle valve. Low denticular plate situated along median one-third of hinge. Ventral process and process pits obscure in material available. Muscle field moderately impressed, triangular in outline, bounded laterally by pair of low, straight ridges which diverge antero-laterally at about 65° to one another. Low, rounded groove longitudinally bisects muscle field which occupies up to one-third length of valve. Individual scars not distinguished.

Area immediately adjacent to muscle field may be fairly coarsely tuberculate.

Interior of brachial valve. Notothyrial platform low, produced anteriorly as low, convex myophragm which bisects muscle field longitudinally. Cardinal process lobes pear-shaped in outline, separated by shallow sub-angular groove which may bear faint ridge along its mid-length. Socket ridges short, immediately lateral to cardinal process lobes and antero-laterally divergent at about 140° to one another.

Muscle field moderately well impressed, occupying about one-third length of valve and one-fifth width. Small, median pair of adductor scars separated by myophragm and bounded laterally by second, semi-elliptical pair. Muscle field bounded laterally by pair of low ridges.

Dimensions of figured specimens in mm

		Length	Width
Partially exfoliated exterior of pedicle valve and counterpart impression	Holotype, NMW 70.3G.42a, b	13.4	19.8
Artificial (calcined) internal mould of pedicle valve	NMW 70.3G.43a	12.4	16.8 est.
Internal mould of pedicle valve	NMW 70.3G.44	13.5	19.2 est.
Internal mould of brachial valve	NMW 70.3G.45	—	—

Holotype. NMW 70.3G.42a, b, Plate 60, figs. 10a, b.

Distribution. *P. (M.) deflecta* is known at present only from the Usk inlier where it occurs very commonly in the upper part of the Wenlock Limestone and in the siltstones immediately overlying the Wenlock Limestone; these siltstones are regarded by Walmsley (1959, p. 487) as being of Wenlock age.

Pholidostrophia (*Mesopholidostrophia*) *salopiensis* Cocks

Plate 60, figs. 6-9

1967 *Pholidostrophia salopiensis* Cocks, p. 263, pl. 39, figs. 12-17.

Discussion. When Cocks (1967, p. 263) originally described this species he commented that it lacked a nacreous shell and suggested that it should therefore be assigned to an unnamed subgenus of *Pholidostrophia*, based on *Stropheodonta* (*Brachyprion*) *sefinensis* Williams 1951, which was to be described by Harper, Johnson, and Boucot. These authors subsequently (1967, p. 411) made *S. (B.) sefinensis* the type species of *Eopholidostrophia*, a new genus which differs from *P. (Mesopholidostrophia)* in having well-developed radial ornament and an obscure dorsal muscle field. The species *salopiensis* has faint radial ornament and a well-developed dorsal muscle field, which excludes it from *Eopholidostrophia*; it is considered here instead to be an early representative of *P. (Mesopholidostrophia)*. Although Cocks inferred that *salopiensis* should belong to *Eopholidostrophia* since it lacked a nacreous shell lustre, this feature is clearly variable within a species and its presence or absence cannot be regarded as being diagnostic (Harper, Johnson, and Boucot 1967, pp. 415-16). Specimens collected by the writer are in the form of moulds so that it is not possible to comment on the nature of the shell material.

Comparison. *P. (M.) salopiensis* resembles both *P. (M.) laevigata* and *P. (M.) nitens* in size and outline but may be distinguished by its strongly convex pedicle valve.

Description. Exterior. Shells small, concavo-convex with pedicle valve 2-3 times as deep as brachial valve. Outline semicircular, hinge-line straight, produced as short, rounded ears. Margins of shell evenly curved. Commissures smooth, anterior commissure rectimarginate.

Ventral interarea plane, apsacline. Delthyrium appears to be open. Dorsal interarea obscure; notothyrium appears to be partially occupied by cardinal process and closed by chilidium.

Radial ornament weak. Very fine parvicostellae present over whole shell, tending to become better developed anteriorly. Faint concentric growth-lines visible on some specimens.

Interior of pedicle valve. Median one-third of hinge bears pair of low denticular plates. Dental plates absent. Muscle field broadly triangular in outline, occupying about one-third width of valve and about half length. Pair of low, straight or gently curved ridges bound muscle field laterally. Adductor scars narrow, elongated, situated medially and separated longitudinally by low myophragm. Relatively broad, suboval diductor scars enclose adductor scars laterally and antero-laterally.

Interior of brachial valve. Cardinal process lobes small, rounded, supported on very low notothyrial platform. Socket plates short, antero-laterally divergent at about 130° to one another. Muscle field relatively well impressed, confined to posterior quarter of

valve and bisected longitudinally by low, flat ridge originating from notothyrial platform. Adductor scars oval, unbounded either laterally or anteriorly.

Dimensions of figured specimens in mm

		<i>Length</i>	<i>Width</i>
Internal mould of pedicle valve	NMW 70.3G.38	7.7	9.3
Internal mould of pedicle valve	NMW 70.3G.39a	7.3	8.9
Internal mould of pedicle valve and counterpart external mould	NMW 70.3G.40a, b	5.4	7.4
External mould of brachial valve	NMW 70.3G.41a	6.2	8.0

Type specimens. Holotype, GSM 102720 (Cocks 1967, pl. 39, fig. 15) from the Upper Llandovery Purple Shales of Domas, Shropshire. The remainder of the material figured by Cocks is in the Oxford University Museum.

Distribution. *P. (M.) salopiensis* occurs fairly commonly in beds mapped by Williams (1953b, pp. 198–9) as lower Wenlock in age, in the Llandovery–Llandeilo district of Carmarthenshire. It has previously been recorded by Cocks (1967, p. 264) only from beds of Upper Llandovery (C₄₋₆) age in the Welsh Borderland.

Pholidostrophia (Mesopholidostrophia) fletcheri (Davidson)

Plate 60, figs. 14–16

1847 *Leptaena fletcheri* Davidson, p. 58, pl. 12, figs. 9–11.

1848 *Leptaena fletcheri* Davidson; Davidson, p. 319, pl. 3, fig. 12.

1871 *Strophomena fletcheri* (Davidson); Davidson, p. 317, pl. 47, figs. 5, 6.

Comparison. *P. (M.) fletcheri*, which is known only from 3 syntype specimens, differs from all other pholidostrophiids described in this paper by having a more oval outline, with the point of maximum width at about the mid-length of the shell. It is also distinguished by being gently concavo-convex posteriorly and becoming fairly strongly concavo-convex anteriorly.

Description. Exterior. Outline sub-oval to semicircular, gently concavo-convex posteriorly, becoming fairly strongly concavo-convex in anterior half of shell. Hinge-line straight, cardinal angles well rounded. Although the syntypes are broken, point of maximum width appears to be at about mid-length of shell, and hinge width equals about four-fifths of maximum width. Lateral and anterior margins evenly curved. Commissures smooth, anterior commissure rectimarginate.

Ventral interarea plane to very gently concave, weakly apsacline, 2–3 times as long as dorsal interarea, which is anacline. Delthyrium small, with convex, apical pseudo-deltidium; delthyrial angle 70–80°. Notothyrium completely sealed by convex chilidium. Both interareas show traces of fine internal denticulation extending for almost two-thirds width of hinge.

Surface of both valves almost smooth, but very faint radial ornament may be distinguished.

Interiors unknown.

Dimensions in mm		Length	Maximum width	Hinge width
Articulated shell	BM B5821	13.9	—	—
Exterior of brachial valve	BM BB32803 (ex B5821)	10.1	16.8 est.	—
Exterior of brachial valve	BM BB32804 (ex B5821)	—	—	—

Type specimens. 3 specimens in the British Museum bear a label in Davidson's handwriting identifying them as types; none is readily identifiable with Davidson's original (1847, pl. 12, figs. 9–11) or later figures (1848, pl. 3, fig. 12; 1871, pl. 47, figs. 5, 6), but it is almost certain that in all 3 cases the drawings were composite reproductions based on the 3 syntypes. The most complete of these specimens (BM B5821) is selected as lectotype and is figured on Plate 60, figs. 15 a–c. The remaining two syntypes (BM BB32804 and BM BB32803) are figured on Plate 60, figs. 14, 16 respectively.

Distribution. All three syntypes are from the Wenlock Limestone of Benthall Edge, Shropshire.

Acknowledgements. I thank Dr. V. G. Walmsley for his help and supervision during the early stages of this work at University College, Swansea, where research facilities were provided by Professor F. H. T. Rhodes. Dr. L. R. M. Cocks (British Museum, Natural History) has discussed problems of stropheodontid systematics with me on a number of occasions, and both he and Dr. Walmsley read and criticized a first draft of the manuscript.

The following people kindly allowed me to borrow and study specimens in their care: Dr. D. L. Bruton (Paleontologisk Museum, Oslo), Dr. L. R. M. Cocks (British Museum, Natural History), Dr. M. L. K. Curtis (City Museum, Bristol), Dr. V. Jaanusson (Naturhistoriska Riksmuseet, Stockholm), Dr. C. E. O'Riordan (National Museum of Ireland), Dr. R. B. Rickards (Sedgwick Museum, Cambridge), Dr. I. Strachan (Birmingham University), Mr. D. E. White (Institute of Geological Sciences, London), and Mr. M. White (Bristol University Museum).

Financial aid from the Sir Richard Stapley Educational Trust, the Natural Environment Research Council, and the National Museum of Wales is gratefully acknowledged.

REFERENCES

- BANCROFT, B. B. 1949. *Welsh Valentian Brachiopods and the Strophomena Antiquata Group of Fossil Brachiopods* (ed. LAMONT, A.). 16 pp., 3 pl. Privately printed. Mexborough.
- BASSETT, M. G. 1970. The articulate brachiopods from the Wenlock Series of the Welsh Borderland and South Wales. *Palaeontogr. Soc. [Monogr.]*, Pt. 1, 1–26, pl. 1–3.
- BOUCOT, A. J., JOHNSON, J. G., HARPER, C. W., and WALMSLEY, V. G. 1966. Silurian brachiopods and gastropods of southern New Brunswick. *Bull. geol. Surv. Can.* **140**, 1–45, pl. 1–18.
- CASTER, K. E. 1939. A Devonian fauna from Colombia. *Bull. Am. Paleont.* **24** (83), 1–218, pl. 1–14.
- 1945. New names for two homonyms. *J. Paleont.* **19**, 319.
- COCKS, L. R. M. 1967. Llandovery stropheodontids from the Welsh Borderland. *Palaeontology*, **10**, 245–65, pl. 37–39.
- CURTIS, M. L. K. 1955. In CURTIS, M. L. K., DONOVAN, D. T., KELLAWAY, G. A., and WELCH, F. B. A. *Bristol and its Adjoining Counties*, chap. 1, *Geology*, 3–33, pl. 37. Bristol (British Association).
- 1956. Type and figured specimens from the Tortworth inlier, Gloucestershire. *Proc. Bristol Nat. Soc.* **29**, 147–54.
- DALMAN, J. W. 1828. Uppställning och Beskrifning af de i Sverige funne Terebratuliter. *K. Svenska Vetensk. Akad. Handl.* [for 1827], 85–155, pl. 1–6.
- DAVIDSON, T. 1847. Observations on some of the Wenlock-limestone Brachiopoda, with descriptions of several new species. *London Geol. J.* **1**, 52–65, pl. 12, 13.
- 1848. Mémoire sur les brachiopodes du système silurien supérieur d'Angleterre. *Bull. Soc. géol. Fr.* (2), **5**, 309–38, pl. 3, 4 (pars).
- 1866–71. A monograph of the British fossil Brachiopoda, vol. 3, part 7, Silurian. *Palaeontogr.*

- Soc. [Monogr.]: No. 1, 1866, 1-88, pl. 1-12; No. 2, 1867, 89-168, pl. 13-22; No. 3, 1869, 169-248, pl. 23-27; No. 4, 1871, 249-397, pl. 38-50.
- DAVIDSON, T. 1883. A monograph of the British fossil Brachiopoda, vol. 5, part 2, Silurian Supplement. Ibid. 135-242, pl. 8-17.
- DIXON, E. E. L. 1921. The geology of the South Wales Coalfield. Part 13, The country around Pembroke and Tenby. *Mem. geol. Surv. U.K.* i-vi, 220 pp., 5 pl.
- GROOM, T. and LAKE, P. 1908. The Bala and Llandovery rocks of Glyn Ceiriog, North Wales. *Q. Jl geol. Soc. Lond.* 64, 546-95, pl. 53.
- HALL, J. and CLARKE, J. M. 1892-94. An introduction to the study of the genera of Palaeozoic Brachiopoda. *N.Y. State Geol. Survey, Paleont. N.Y.* 8 (1), 1892, i-xvi, 1-367, 41 pl.; (2), 1894, i-xvi, 1-394, pl. 21-84.
- HARPER, C. W., JOHNSON, J. G., and BOUCOT, A. J. 1967. The Pholidostrophiinae (Brachiopoda; Ordovician, Silurian, Devonian). *Senckenberg. leth.* 48, 403-61, pl. 1-10.
- HAVLÍČEK, V. 1967. Brachiopoda of the Suborder Strophomenidina in Czechoslovakia. *Rozpr. ústřed. Úst. geol.* 33, 1-235, pl. 1-52.
- HEDE, J. E. 1960. In REGNELL, G. and HEDE, J. E. The lower Palaeozoic of Scania—the Silurian of Gotland. Guide to excursions A22 and C17. *Rept. Int. geol. Congr.*, 21st Sess., Norden, 1-87.
- HISINGER, W. 1828. *Anteckningar i Physik och Geognosi under resor uti Sverige och Norrige*. Haft 4, 1-258, pl. 1-9. Stockholm.
- HOLLAND, C. H., LAWSON, J. D., and WALMSLEY, V. G. 1963. The Silurian rocks of the Ludlow district, Shropshire. *Bull. Br. Mus. nat. Hist. (Geol.)*, 8, 93-171, pl. 1-7.
- HOLTEDAHN, O. 1916. The Strophomenidae of the Kristiania region. *K. norske Vidensk. Selsk. Skr.* 12, 1-117, pl. 1-16.
- LAWSON, J. D. 1960. The succession of shelly faunas in the British Ludlovian. *Rept. Int. geol. Congr.*, 21st Sess., Norden, 7, 114-25.
- LINDSTRÖM, G. 1861. Bidrag till Kännedomen om Gotlands Brachiopoder. *Öfvers. K. Vetensk. Akad. Förh.* 17 [for 1860], 337-82, pl. 12, 13.
- MCCOY, F. 1846. *A Synopsis of the Silurian Fossils of Ireland*, 72 pp., 5 pl. Dublin.
- 1851. On some new Cambro-Silurian Fossils. *Ann. Mag. nat. Hist.* 8, 387-409.
- 1851-55. In SEDGWICK, A. and MCCOY, F. *A Synopsis of the Classification of the British Palaeozoic Rocks, with a Systematic Description of the British Palaeozoic Fossils in the Geological Museum of the University of Cambridge*. (1), 1851, i-iv, 1-184; (2), 1852, i-viii, 185-406; (3), 1855, i-xcviii, 407-661. 25 pl. London and Cambridge.
- MURCHISON, R. I. 1839. *The Silurian System*, i-xxxii, 768 pp., 37 pl. London.
- 1854. *Siluria*. i-xvi, 1-523, pl. 1-37.
- PANDER, C. H. 1830. *Beiträge zur Geognosie des Russischen Reiches*. 165 pp., 31 pl. St. Petersburg.
- PHILLIPS, J. and SALTER, J. W. 1848. Palaeontological appendix to Professor John Phillips' Memoir on the Malvern Hills compared with the Palaeozoic districts of Abberley, etc. *Mem. geol. Surv. U.K.* 2 (1), 331-86, pl. 4-30.
- SALTER, J. W. 1873. *A Catalogue of the Collection of Cambrian and Silurian Fossils contained in the Geological Museum of the University of Cambridge*, i-xlvi, 204 pp. Cambridge.
- SHALER, N. S. 1865. List of the Brachiopoda from the Island of Anticosti, sent by the Museum of Comparative Zoology to different Institutions in Exchange for other Specimens, with Annotations. *Bull. Mus. comp. Zool. Harv.* 1 (4), 61-70.
- WALMSLEY, V. G. 1959. The geology of the Usk Inlier (Monmouthshire). *Q. Jl geol. Soc. Lond.* 114 [for 1958], 483-521, pl. 22.
- WHITTARD, W. F. 1932. The stratigraphy of the Valentian rocks of Shropshire. The Longmynd-Shelve and Breidden outcrops. Ibid. 87, 859-902, pl. 58-62.
- (ed.). 1961. *Lexique Stratigraphique International*, 1, Europe. Fasc. 3a, England, Wales and Scotland. Part 3aV, Silurian. 273 pp. Paris.
- WILLIAMS, A. 1950. New stropheodontid brachiopods. *J. Wash. Acad. Sci.* 40, 277-82, 1 pl.
- 1951. Llandovery brachiopods from Wales with special reference to the Llandovery district. *Q. Jl geol. Soc. Lond.* 107, 85-136, pl. 3-7.
- 1953a. North American and European stropheodontids: their morphology and systematics. *Mem. geol. Soc. Am.* 56, 1-67, pl. 1-13.

- WILLIAMS, A. 1953*b*. The geology of the Llandeilo District, Carmarthenshire. *Q. Jl geol. Soc. Lond.* **108** [for 1952], 177–208, pl. 9.
- 1962. The Barr and Lower Ardmillan Series (Caradoc) of the Girvan district, south-west Ayrshire, with descriptions of the Brachiopoda. *Mem. geol. Soc. Lond.* **3**, 1–267, pl. 1–25.
- 1965. In MOORE, R. C. (ed.). *Treatise on Invertebrate Paleontology, Part H, Brachiopoda*. 2 vols., i–xxxii, 927 pp., 746 figs. Geol. Soc. Amer. and Univ. Kansas Press.
- WOODS, H. 1891. *Catalogue of the Type Fossils in the Woodwardian Museum, Cambridge*. i–xvi, 180 pp. Cambridge.

MICHAEL G. BASSETT
Department of Geology
National Museum of Wales
Cardiff, CF1 3NP

Typescript received 3 August 1970