

HOMOTRYPA AND AMPLEXOPORA? FROM THE CARADOC SERIES, SHROPSHIRE

by JUNE R. PHILLIPS ROSS

ABSTRACT. Additional bryozoan species from the *Harknessella subquadrata* horizon of the Hoar Edge Limestone, Evenwood Quarry, include unique inverted cone-shaped colonies of *Homotrypa oweni* sp. nov. and two species questionably assigned to *Amplexopora*, *Amplexopora? evenensis* sp. nov. and *Amplexopora? sp. A*.

MATERIAL collected in 1963 by Dr. D. Owen, Manchester University Museum, from the upper part of the Evenwood Quarry, Shropshire, in the *Harknessella subquadrata* horizon of the Hoar Edge Limestone, permits observations on bryozoan species of *Homotrypa* and *Amplexopora?* that add to the bryozoan fauna described by Ross (1963). This additional material in a large slab from Evenwood Quarry includes a great abundance of *Homotrypa oweni* sp. nov. Associated with it are *Phaenopora stubblefieldi* Ross and *Amplexopora? evenensis* sp. nov., which are common, and fragments of *Amplexopora thomasi* Ross and *Amplexopora? sp. A*, which are sparse.

The three newly discovered species, *Homotrypa oweni*, *Amplexopora? evenensis*, and *Amplexopora? sp. A*, are distinctly different from previously described species of these genera. The distinctive inverted-cone shape of colonies of *Homotrypa oweni* permits ready identification of this species. However, the phylogenetic affinities of this species are at present not determinable as it appears to have little similarity with previously described species. *Amplexopora? evenensis* and *Amplexopora? sp. A* both display strongly crenulate zooecial walls which thicken only slightly in the narrow peripheral regions. Such a marked crenulation of the zooecial walls as is seen in these two species has not been noted so far in other species of the genus *Amplexopora*, so that these two Caradocian species have been assigned with question to *Amplexopora*. The concentration of diaphragms in the subperipheral region and the narrowness of the peripheral region are distinctive features of *Amplexopora? evenensis*. *Amplexopora? sp. A* has isolate cystiphragms on its distal walls, which together with limited diaphragms in the zooecial tubes and numerous diaphragms in the mesopores, single out this species. *Amplexopora? evenensis* has some general similarities in the structure of the peripheral region and size and shape of acanthopores with certain species of *Amplexopora* from the Champlainian and Cincinnati Series of North America, including such species as *A. winchelli* (Ulrich), *A. pustulosa* Ulrich, *A. ampla* Ulrich and Bassler, and *A. convoluta* Bassler; however, for reasons noted above it has no strong affinities with any of these species. *Amplexopora? sp. A* has little similarity to previously described species.

SYSTEMATIC DESCRIPTIONS

Family MONTICULIPORIDAE

Genus HOMOTRYPA Ulrich

Homotrypa oweni sp. nov.

Plate 2, figs. 1-5; Plate 3, fig. 6; Table 1

[Palaeontology, Vol. 8, Part 1, 1965, pp. 5-10, pl. 2-3.]

Material. Holotype, LL 2807A; three figured paratypes, LL 2807B-D; 8 unfigured thin-sectioned paratypes, LL 2807E-L; three unfigured hand-specimen paratypes, LL 2807M-O.

Description. Hollow, inverted-cone shaped colonies encrust calcite fragments and crinoid columnals (Pl. 2, fig. 5). The cones are about 2 to 3 mm. in diameter at the tapered end and expand distally to 4 mm.

The subpolygonal zooecial openings interlock in a regular arrangement and cystiphragms are visible across the zooecial openings (Pl. 2, figs. 1, 2, 5). Short zooecial tubes are reclined for a short distance in the proximal part along the basal lamina; zooecial tubes are straight for the remainder of their length (Pl. 2, figs. 3-5; Pl. 3,

TABLE 1. Measurements of *Homotrypa oweni* sp. nov. (in millimetres)

Catalogue no.	LL 2807B	LL 2807A	LL 2807C
Diameter of colony	Not determined	2.6 to 3.8	2.6 to 4.1
Length of colony	Not determined	12.5	9
No. of zooecia per 2 mm. longitudinally	9 to 10	8 to 10	9 to 11
Diameter of zooecial opening, max.	0.24 × 0.24	0.24 × 0.22	0.18 × 0.18
min.	0.13 × 0.16	0.18 × 0.18	0.15 × 0.13
Combined thickness of adjacent zooecial walls in peripheral region	0.01 to 0.02	0.01 to 0.02	0.02 to 0.03
Diameter of mesopore, max.	0.08 × 0.06	0.08 × 0.10	0.08 × 0.10
min.	0.04 × 0.04	0.05 × 0.05	0.03 × 0.03
Diameter of acanthopore	0.02	0.01 to 0.02	0.02
No. of acanthopores per zooecium	1 to 3	2 to 3	1 to 4
No. of diaphragms per 1 mm. in zooecial tube in peripheral region	Not determined	1 to 2	Not determined

fig. 6). Diaphragms are sparse in the zooecial tubes, commonly one diaphragm in the upper part or reclined portion of zooecial tube and another diaphragm more distally located; diaphragms more closely spaced in mesopores, 4 to 5 in length of mesopore. Mesopores develop in distal region of reclined portion of zooecial tubes (Pl. 2, fig. 5). Cystiphragms occur almost always as regular overlapping structures on the distal walls of zooecial tubes and rarely occur on the proximal zooecial walls; they develop just above the inclined portion of zooecial tubes and succeeding cystiphragms overlap by a half to one-third the preceding cystiphragms (Pl. 2, figs. 3-5; Pl. 3, fig. 6). Acanthopores are small and indistinct and are difficult to locate in longitudinal sections; in tangential sections they occur at the junctions of zooecial and mesopore walls and appear as small dark spots (Pl. 2, fig. 2). Two to four mesopores occur around each zooecium but do not isolate zooecia. Zooecial walls thin; they display laminate wall structure in which the laminae are steeply inclined along the inner part of the zoo-

EXPLANATION OF PLATE 2

Figs. 1-5. *Homotrypa oweni* sp. nov. 1, Deep tangential section across more proximal and medial parts of colony, paratype LL 2807C, × 20. 2, Tangential section showing thin-walled zooecial tubes, small acanthopores, and cystiphragms across zooecial openings, paratype LL 2807B, × 50. 3, 4, Parts of longitudinal sections showing basal lamina, reclined proximal portions of zooecial tubes, and numerous cystiphragms and sparse diaphragms in more distal portions of zooecial tubes, holotype LL 2807A, paratype LL 2807D, × 50. 5, Oblique longitudinal section through encrusting tapering colony, holotype LL 2807A, × 20.

zoecial tube and curve for a short distance in the outer part of zoecial wall where they abut against laminae of adjacent zoecial walls.

Remarks. In this species the form of the colony is most distinctive; both the cone shape and the encrusting habit. This species is not closely comparable to any previously described species of *Homotrypa* and in internal features is characterized by regular, overlapping cystiphragms on the distal zoecial walls and sparse diaphragms in the zoecial tubes, indistinct acanthopores, and an abundance of mesopores which are evenly spaced between zoecial openings.

This species is also present in collection RR 2669 in the Geological Survey Museum, London, from Wylde's Quarry, 30 chains south-west of Harnage Grange, Shropshire (Pocock *et al.* 1938, fig. 28). The species is named after Dr. D. Owen, Director, Manchester University Museum.

Family AMPLEXOPORIDAE Miller
Genus AMPLEXOPORA Ulrich

Amplexopora? evenensis sp. nov.

Plate 3, figs. 1-3; Table 2

Material. Holotype, LL 2823A. Paratypes, LL 2823B figured, LL 2823C-L unfigured.

Description. Robust cylindrical to subglobular branching colonies; stems 4 to 5 mm. in diameter; some small overgrowths. Polygonal zoecial openings are enclosed by very slender amalgamate zoecial walls (Pl. 3, fig. 1). Polygonal mesopores are interspersed between zoecial tubes and small, though at times indistinct, acanthopores commonly penetrate the junctions of the zoecial walls and, in some instances, a second set of acanthopores are located near the junctions of the zoecial walls and may lie near the inner part of the zoecial walls. Long zoecial tubes with broadly crenulate zoecial walls slope steeply from the axial region and pass into the very narrow peripheral region where zoecial walls are closely crenulated (Pl. 3, figs. 2, 3). Diaphragms are absent in the axial region, closely spaced in the subperipheral region, and absent in the peripheral region. Diaphragms in the subperipheral region may be flat, curved, or overlapping cystoidal. Acanthopores, commonly difficult to observe in longitudinal sections, appear as narrow rods with walls of steeply inclined laminae; some extend from the subperipheral region and others project above the zoarial surface. Deep tangential sections display only a few acanthopores and expanding mesopores. Zoecial walls are slightly thickened in the peripheral region, where they display inclined laminae in the inner part of the walls. These laminae are curved in the outer part of the zoecial walls and are amalgamate in a dark irregular band with laminae of the adjacent wall. Mesopores arise in the subperipheral region, extend to the periphery, and have sparse diaphragms in the subperipheral region. At three to four corresponding levels in the colonies, there is bifurcation of the zoecial tubes with a marked increase in the diameter of the zoarium; sometimes associated with this bifurcation of the tubes is a curved band of one or two diaphragms which extends across the colony.

Remarks. This species is characterized by slender, crenulate zoecial walls, small but

distinct acanthopores, some of which extend into the subperipheral region, and a small number of diaphragms in the subperipheral region of the zooecial tubes and mesopores. *Amplexopora? evenensis* has little similarity with previously described species, which generally display thicker walls, larger zooecial tubes, and a greater abundance of diaphragms. *Amplexopora winchelli* (Ulrich) (1893, 'Middle third of Trenton shales, at St. Paul', pl. 27, figs. 1, 4, 6) has polygonal zooecial openings, small acanthopores, diaphragms in the subperipheral region, and a lack of diaphragms in the peripheral region. However, Ulrich's other illustrations (pl. 27, figs. 2, 3, 5, 7, 8) show thicker-walled zooecia, more numerous and larger acanthopores, and a great abundance of

TABLE 2 Measurements of *Amplexopora? evenensis* sp. nov. (in millimetres)

Catalogue no.	LL 2823B	LL 2823A
Diameter of colony	Not determined	4
No. of zooecia per 2 mm.	9 to 11	9 to 11
Diameter of zooecial opening, max.	0.27 × 0.28	0.29
min.	0.16 × 0.20	0.17
Combined thickness of adjacent zooecial walls in peripheral region	0.02	0.01 to 0.03
Diameter of mesopore, max.	0.16 × 0.05	Not determined
min.	0.05 × 0.02	0.04
Diameter of acanthopore	0.01 to 0.03	0.01 to 0.02
No. of acanthopores per zooecium	2 to 5	Not determined
No. of diaphragms per 1 mm. in zooecial tube in subperipheral region	Not determined	3 to 4

diaphragms. *A. pustulosa* Ulrich (1890, 'Cincinnati group, Hanover, Clarksville, and other localities in Ohio') likewise, while displaying some similarities in polygonal zooecial openings and small acanthopores, is considerably more robust in its zooecial and zoarial structures. *A. ampla* Ulrich and Bassler (1904, 'Fairmount beds, Cincinnati, Ohio') has larger zooecial tubes and diaphragms in the axial region. Slender zooecial walls, narrow peripheral region, and small acanthopores suggest similarities with *A.? evenensis*.

This new species takes its name from Evenwood Quarry.

EXPLANATION OF PLATE 3

- Figs. 1-3. *Amplexopora? evenensis* sp. nov. 1, Deep tangential section showing polygonal zooecial openings and mesopores, and small acanthopores at junctions of zooecial walls, paratype LL 2823B, × 50. 2, Peripheral region of part of longitudinal section showing thickened crenulate zooecial walls, holotype LL 2823A, × 50. 3, Part of longitudinal section showing sparse diaphragms in axial region and concentration of diaphragms in subperipheral region, holotype LL 2823A, × 20.
- Figs. 4, 5, 7, 8. *Amplexopora? sp. A.* 4, Longitudinal section showing crenulate zooecial walls and sparse diaphragms in zooecial tubes and mesopores and occasional cystiphragm in zooecial tube, LL 2824B, × 50. 5, Longitudinal section showing general aspect of zooecia and mesopores, LL 2824B, × 20. 7, Deep tangential section showing round to subpolygonal zooecial openings, polygonal mesopores, and an occasional acanthopore at junction of zooecial wall, LL 2824A, × 50. 8, Longitudinal section showing part of overgrowth with very distinct cystiphragms, LL 2824B, × 50.
- Fig. 6. *Homotrypa oweni* sp. nov. Part of longitudinal section showing laminate structure of zooecial walls and cystiphragms in zooecial tubes, paratype LL 2807D, × 100.

Amplexopora? sp. A

Plate 3, figs. 4, 5, 7, 8; Table 3

Material. Figured specimens, LL 2824A, 2824B.

Description. Very slender cylindrical branches with overgrowths (Pl. 3, figs. 5, 8). Round to subpolygonal zooecial openings are enclosed by slender amalgamate zooecial walls (Pl. 3, fig. 7). Mesopores and occasionally acanthopores occur at junctions of zooecial walls. Long slender zooecial tubes with slightly crenulate walls slope steeply from the axial region and curve abruptly into subperipheral region (Pl. 3, fig. 5). Narrow peripheral region displays limited thickening of crenulate zooecial walls and extension of

TABLE 3. Measurements of *Amplexopora?* sp. A (in millimetres)

<i>Catalogue no.</i>	<i>LL 2824A</i>	<i>LL 2824B</i>
Diameter of colony	Not determined	1.6
No. of zooecia per 2 mm. longitudinally	10 to 11	11 to 13
Diameter of zooecial opening, max.	0.18 × 0.16	0.15
min.	0.10 × 0.13	0.09
Combined thickness of adjacent zooecial walls in peripheral region	0.05 to 0.08	0.01 to 0.02
Diameter of mesopore, max.	0.08 × 0.10	0.11
min.	0.02 × 0.02	0.04
Diameter of acanthopore	0.01 to 0.02	0.02
No. of acanthopores per zooecium	2 to 3	Not determined
No. of diaphragms in zooecial tube in peripheral region	Not determined	4 in 0.3 mm.

mesopores from subperipheral region. Occasionally one or two diaphragms are present in subperipheral region or peripheral region of zooecial tube. Diaphragms are abundant and closely spaced in mesopores (Pl. 3, figs. 4, 5). An occasional isolate or overlapping set of two or three cystiphragms occur on distal walls of some zooecial tubes (Pl. 3, figs. 4, 5). Acanthopores extending from subperipheral region to the periphery are almost as wide as the zooecial walls but are not readily observable in longitudinal section. The slightly thickened zooecial walls of the peripheral region have a wall structure as in *Amplexopora?* *evenensis*.

Remarks. *Amplexopora?* sp. A has certain similarities to *A.?* *evenensis* in the arrangement of the zooecial tubes, crenulate zooecial walls, laminate zooecial wall structure, and arrangement of the zooecial openings. It differs from *A.?* *evenensis* in having more slender colonies, narrower zooecial tubes, very few diaphragms in the zooecial tubes, cystiphragms in the zooecial tubes, thicker zooecial walls in the peripheral region, and an abundance of diaphragms in the mesopores. The species has little similarity to previously described species of *Amplexopora* and its assignment to the genus is questioned.

Acknowledgements. I am most grateful to Dr. D. E. Owen, Director, Manchester Museum, for kindly sending this material to me for examination and study.


Repository. All the specimens referred to are in the Manchester Museum, The University, Manchester 13.

REFERENCES


- POCOCK, R. W., WHITEHEAD, T. H., WEDD, C. B., and ROBERTSON, T. 1938. Shrewsbury District including the Hanwood Coalfield. *Mem. geol. Surv. U.K.*
- ROSS, J. R. P. 1963. Trepostome Bryozoa from Caradoc Series, Shropshire. *Palaontology*, **6**, 1-11, pl. 1-3.
- ULRICH, E. O. 1890. Palaeozoic Bryozoa. *Illinois geol. Surv., Geology and Palaeontology*, **8**, 283-688, pl. 29-78.
- 1893. On Lower Silurian Bryozoa of Minnesota. *Minnesota geol. and nat. Hist. Surv., Geology of Minnesota*, **3** (1), *Paleontology*, 96-332, 28 pl., 1895. (Author's separate, 1893).
- and BASSLER, R. S. 1904. A revision of the Paleozoic Bryozoa. Pt. 2, Trepostomata. *Smithson. misc. Coll.* **47**, 15-55, pl. 6-14.

JUNE R. PHILLIPS ROSS
Department of Geology,
University of Illinois,
Urbana, Illinois, U.S.A.

Manuscript received 13 January 1964



PHILLIPS ROSS, Ordovician Bryozoa



PHILLIPS ROSS, Ordovician Bryozoa